

1. RINGS AND MODULES

Recall that a *ring* is a system $(R, +, \cdot, 0, 1)$ consisting of a set R , two binary operations, addition $(+)$ and multiplication (\cdot) , and two elements $0 \neq 1$ of R , such that $(R, +, 0)$ is an abelian group, $(R, \cdot, 1)$ is a monoid (i.e., a semigroup with identity 1) and multiplication is left and right distributive over addition. A ring whose multiplicative structure is abelian is called a *commutative ring*.

Example 1.1. (1) $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$ are commutative rings.

- (2) Let K be a field; the ring $K[x_1, \dots, x_n]$ of polynomials in the indeterminates x_1, \dots, x_n is a commutative ring.
- (3) Let K be a field; consider the ring $R = M_n(K)$ of $n \times n$ -matrices with coefficients in K with the usual "rows times columns" product. Then R is a non-commutative ring

Definition 1.2. A left R -module is an abelian group M together with a map $R \times M \rightarrow M$, $(r, m) \mapsto rm$, such that for any $r, s \in R$ and any $x, y \in M$

- M1 $r(x + y) = rx + ry$
 M2 $(r + s)x = rx + sx$
 M3 $(rs)x = r(sx)$
 M4 $1x = x$

We write ${}_R M$ to indicate that M is a left R -module.

Example 1.3. (1) Any abelian group G is a left \mathbb{Z} -module by defining, for any $x \in G$ and $n > 0$, $nx = \underbrace{x + \dots + x}_{n \text{ times}}$.

- (2) Given a field K , any vector space V over K is a left K -module.
- (3) Let R be the matrix ring $M_n(K)$ and consider the vector space $V = K^n$. Given a matrix A and a vector $v \in V$, let Av be the usual "rows times columns" product. Then V is a left R -module.
- (4) Any ring R is a left R -module, by using the left multiplication of R on itself. It is called the *regular module*.
- (5) Consider the zero element of the ring R . Then the abelian group $\{0\}$ is trivially a left R -module.

Remark 1.4. Consider M an abelian group and $\text{End}^l(M)$ the ring of the endomorphism of M acting on the left (i.e. $fg(x) = f(g(x))$). A *representation* of R in $\text{End}^l(M)$ is a homomorphism of ring

$$\lambda : R \rightarrow \text{End}^l(M), \quad r \mapsto \lambda(r)$$

From the properties of ring homomorphisms it follows that for any $r, s \in R$ and $x, y \in M$

- (1) $\lambda(r)(x + y) = \lambda(r)x + \lambda(r)y$
 (2) $\lambda(r + s)x = \lambda(r)x + \lambda(s)x$
 (3) $\lambda(rs)x = \lambda(r)(\lambda(s)x)$
 (4) $\lambda(1)x = x$

In other words, we can consider $\lambda(r)$ acting on the elements of M as a left multiplication by the element $r \in R$: then we can define $rx := \lambda(r)x$, and this gives a structure of left R -module on M . Conversely, to any left R -module M , we can associate a representation of R in $\text{End}^l(M)$, by defining $\lambda(r) := rx$.

Similarly, we define right R -modules:

Definition 1.5. A right R -module is an abelian group M together with a map $M \times R \rightarrow M$, $(m, r) \mapsto mr$, such that for any $r, s \in R$ and any $x, y \in M$

- M1 $(x + y)r = xr + yr$
 M2 $x(r + s) = xr + xs$
 M3 $x(rs) = (xr)s$
 M4 $x1 = x$

We write M_R to indicate that M is a right R -module.

For the connection between right modules and representations see Exercise 4.5.

If R is a commutative ring, then left R -modules and right R -modules coincide. Indeed, given a left R -module M with the map $R \times M \rightarrow M$ $(r, m) \mapsto rm$, we can define a map $M \times R \rightarrow M$ $(m, r) \mapsto mr := rm$. This map satisfies the axioms of Definition 1.5 (Verify!) and so M is also a right R -module. The crucial point is that, in the third axiom, since R is commutative we have $x(rs) = (rs)x = (sr)x = s(rx) = (rx)s = (xr)s$.

Example 1.6. Consider the ring $R = M_n(K)$ and V the vector space of the columns $M_{n \times 1}(K)$. This is in an obvious way a left R -module but not a right R -module. Similarly, the vector space of the rows $M_{1 \times n}(K)$ is a right R -module but not a left R -module.

Exercise 1.7. Show that given ${}_R M$, for any $x \in M$ and $r \in R$, we have

- (1) $r0 = 0$
- (2) $0x = 0$
- (3) $r(-x) = (-r)x = -(rx)$

Definition 1.8. Let ${}_R M$ be a left R -module. A subset L of M is a submodule of M if L is a subgroup of M and $rx \in L$ for any $r \in R$ and $x \in L$ (i.e. L is a left R -module under operations inherited from M). We write $L \leq M$.

Example 1.9.

- (1) Let G be a \mathbb{Z} -module. The submodules of G are exactly the subgroups of G .
- (2) Let K a field and V a K -module. The submodules of V are exactly the vector subspaces of V .
- (3) Let R a ring. The submodules of the left R -module ${}_R R$ are the left ideals of R . The submodules of the right R -module R_R are the right ideals of R .

Definition 1.10. Let ${}_R M$ be a left R -module and $L \leq M$. The quotient module M/L is the quotient abelian group together with the map $R \times M/L \rightarrow M/L$ given by $(r, \bar{x}) \mapsto \overline{rx}$.

Remark 1.11. The map $R \times M/L \rightarrow M/L$ given by $(r, \bar{x}) \mapsto \overline{rx}$ is well-defined, since if $\bar{x} = \bar{y}$ then $x - y \in L$ and hence $r(x - y) = rx - ry \in L$, that is $\overline{rx} = \overline{ry}$.

In this part of the course we mainly deal with left modules. So, in the following, unless otherwise is stated, with *module* we always mean *left module*.

2. HOMOMORPHISMS OF MODULES

Definition 2.1. Let ${}_R M$ and ${}_R N$ be R -modules. A map $f : M \rightarrow N$ is a homomorphism if $f(rx + sy) = rf(x) + sf(y)$ for any $x, y \in M$ and $r, s \in R$.

Remark 2.2.

- (1) From the definition it follows that $f(0) = 0$.
- (2) Clearly if f and g are homomorphisms from M to N , also $f + g$ is a homomorphism. Since the zero map is obviously a homomorphism, the set $\text{Hom}_R(M, N) = \{f | f : M \rightarrow N \text{ is a homomorphism}\}$ is an abelian group.
- (3) If $f : M \rightarrow N$ and $g : N \rightarrow L$ are homomorphisms, then $gf : M \rightarrow L$ is a homomorphism. Thus the abelian group $\text{End}_R(M) = \{f | f : M \rightarrow M \text{ is a homomorphism}\}$ has a natural structure of ring, called the *ring of endomorphisms of M* . The identity homomorphism $\text{id}_M : M \rightarrow M$, $m \mapsto m$, is the unity of the ring.

Definition 2.3. Given a homomorphism $f \in \text{Hom}_R(M, N)$, the kernel of f is the set $\text{Ker } f = \{x \in M | f(x) = 0\}$. The image of f is the set $\text{Im } f = \{y \in N | y = f(x) \text{ for } x \in M\}$.

It is easy to verify that $\text{Ker } f \leq M$ and $\text{Im } f \leq N$. Thus we can define the *cokernel* of f as the quotient module $\text{Coker } f = N / \text{Im } f$.

A homomorphism $f \in \text{Hom}_R(M, N)$ is called a *monomorphism* if $\text{Ker } f = 0$. f is called an *epimorphism* if $\text{Im } f = N$. f is called *isomorphism* if it is both a monomorphism and an epimorphism. If f is an isomorphism we write $M \cong N$.

Remark 2.4. (1) For any submodule $L \leq M$ there is a canonical monomorphism $i : L \rightarrow M$, which is the usual inclusion, and a canonical epimorphism $p : M \rightarrow M/L$ which is the usual quotient map.

- (2) For any M the trivial map $0 \rightarrow M$, $0 \mapsto 0$, is a mono. The trivial map $M \rightarrow 0$, $m \mapsto 0$, is an epi.

- (3) The monomorphisms, the epimorphisms and the isomorphisms are exactly the injective, surjective and bijective homomorphisms.

Exercise 2.5. Show that $f \in \text{Hom}_R(M, N)$ is an isomorphism if and only if there exist $g \in \text{Hom}_R(N, M)$ such that $gf = \text{id}_M$ and $fg = \text{id}_N$. In such a case g is unique. (We usually denote g as f^{-1}).

Proposition 2.6. Any $f \in \text{Hom}_R(M, N)$ induces an isomorphism $M/\text{Ker } f \cong \text{Im } f$.

Proof. The induced map $M/\text{Ker } f \rightarrow \text{Im } f$, $\bar{m} \mapsto f(m)$ is a homomorphism. Moreover it is clearly a mono and an epi. \square

The usual homomorphism theorems which hold for groups hold also for homomorphisms of modules.

Proposition 2.7. (1) If $L \leq N \leq M$, then $(M/L)/(N/L) \cong M/L$.

(2) If $L, N \leq M$, denote by $L + N = \{m \in M \mid m = l + n \text{ for } l \in L \text{ and } n \in N\}$. Then $L + N$ is a submodule of M and $(L + N)/N \cong N/(N \cap L)$.

Exercise 2.8. Prove the previous Proposition.

3. EXACT SEQUENCES

Definition 3.1. A sequence of homomorphisms of R -modules

$$\dots \rightarrow M_{i-1} \xrightarrow{f_{i-1}} M_i \xrightarrow{f_i} M_{i+1} \xrightarrow{f_{i+1}} \dots$$

is called exact if $\text{Ker } f_i = \text{Im } f_{i-1}$ for any i .

An exact sequence of the form $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ is called a short exact sequence

Observe that if $L \leq M$, then the sequence $0 \rightarrow L \xrightarrow{i} M \xrightarrow{p} M/L \rightarrow 0$, where i and p are the canonical inclusion and quotient homomorphisms, is short exact (Verify!) Conversely, if $0 \rightarrow M_1 \xrightarrow{f} M_2 \xrightarrow{g} M_3 \rightarrow 0$ is a short exact sequence, then f is a mono, g is an epi, and $M_3 \cong \text{Coker } f$ (Verify!).

The following result is very useful:

Proposition 3.2. Consider the commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & L & \xrightarrow{f} & M & \xrightarrow{g} & N & \longrightarrow & 0 \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \\ 0 & \longrightarrow & L' & \xrightarrow{f'} & M' & \xrightarrow{g'} & N' & \longrightarrow & 0 \end{array}$$

If α and γ are monomorphisms (epimorphisms, or isomorphisms, respectively), then so is β

Proof. (1) Suppose α and γ are monomorphisms and let m such that $\beta(m) = 0$. Then $\gamma(g(m)) = 0$ and so $m \in \text{Ker } g = \text{Im } f$. Hence $m = f(l)$, $l \in L$ and $\beta(m) = \beta(f(l)) = f'(\alpha(l)) = 0$. Since f' and α are mono, we conclude $l = 0$ and so $m = 0$.

- (2) Suppose α and γ are epimorphisms and let $m' \in M'$. Then $g'(m') = \gamma(g(m))$, so $g'(m') = g(\beta(m))$; hence $m' - \beta(m) \in \text{Ker } g' = \text{Im } f'$ and so $m' - \beta(m) = f'(l')$, $l' \in L'$. Let $l \in L$ such that $l' = \alpha(l)$: then $m' - \beta(m) = f'(\alpha(l)) = \beta(f(l))$ and so we conclude $m' = \beta(m - f(l))$. \square

4. EXERCISES

Exercise 4.1. Let ${}_R M$ be a R -module and ${}_R R$ the regular module. Consider the abelian group $\text{Hom}_R(R, M)$ and the map $\varphi : \text{Hom}_R(R, M) \rightarrow M$, $f \mapsto f(1)$. Verify that φ is an isomorphism of \mathbb{Z} -modules.

Exercise 4.2. Let ${}_R M$ and define $\text{Ann}_R(M) = \{r \in R \mid rm = 0 \text{ for any } m \in M\}$. M is called faithful if $\text{Ann}_R(M) = 0$. Verify that $\text{Ann}_R(M)$ is an ideal of R . Verify that M has a natural structure of $R/\text{Ann}_R(M)$ -module, given by the map $R/\text{Ann}_R(M) \times M \rightarrow M$, $(\bar{r}, m) \mapsto rm$. Verify that M over $R/\text{Ann}_R(M)$ is a faithful module.

Exercise 4.3. Let f be a homomorphism of R -modules.
Show that f is a mono if and only if $fg = 0$ implies $g = 0$.
Show f is an epi if and only if $gf = 0$ implies $g = 0$

Exercise 4.4. Consider the ring $R = \begin{pmatrix} K & K \\ 0 & K \end{pmatrix}$. Show that $P_1 = \left\{ \begin{pmatrix} k & 0 \\ 0 & 0 \end{pmatrix} \mid k \in K \right\}$ and $P_2 = \left\{ \begin{pmatrix} 0 & k_1 \\ 0 & k_2 \end{pmatrix} \mid k_1, k_2 \in K \right\}$ are left submodules of ${}_R R$. Show that $Q_1 = \left\{ \begin{pmatrix} k_1 & k_2 \\ 0 & 0 \end{pmatrix} \mid k_1, k_2 \in K \right\}$ and $Q_2 = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & k \end{pmatrix} \mid k \in K \right\}$ are right submodules of R_R

Exercise 4.5. Consider M an abelian group and $\text{End}^r(M)$ the ring of the endomorphism of M acting on the right (i.e. $(x)fg = ((x)f)g$). Show that any representation of R in $\text{End}^r(M)$ corresponds to a right R -module M_R .

5. SUMS AND PRODUCTS OF MODULES

Let I be a set and $\{M_i\}_{i \in I}$ a family of R -modules. The cartesian product $\prod_I M_i = \{(x_i) | x_i \in M_i\}$ has a natural structure of left R -module, by defining the operations component-wise:

$$(x_i)_{i \in I} + (y_i)_{i \in I} = (x_i + y_i)_{i \in I}, \quad r(x_i)_{i \in I} = (rx_i)_{i \in I}.$$

This module is called the *direct product* of the modules M_i . It contains a submodule

$$\bigoplus_I M_i = \{(x_i) | x_i \in M_i \text{ and } x_i = 0 \text{ for almost all } i \in I\}$$

Recall that "almost all" means "except for a finite number". The module $\bigoplus_I M_i$ is called the *direct sum* of the modules M_i . Clearly if I is a finite set then $\prod_I M_i = \{(x_i) | x_i \in M_i\} = \bigoplus_I M_i$. For any component $j \in I$ there are canonical homomorphisms

$$\prod_I M_i \rightarrow M_j, \quad (x_i)_{i \in I} \mapsto x_j \quad \text{and} \quad M_j \rightarrow \prod_I M_i, \quad x_j \mapsto (0, 0, \dots, x_j, 0, \dots, 0)$$

called the *projection* on the j^{th} -component and the *injection* of the j^{th} -component. They are epimorphisms and monomorphism, respectively, for any $j \in I$. The same is true for $\bigoplus_I M_i$.

When $M_i = M$ for any $i \in I$, we use the following notations

$$\prod_I M_i = M^I, \quad \bigoplus_I M_i = M^{(I)}, \quad \text{and if } I = \{1, \dots, n\}, \quad \bigoplus_I M_i = M^n$$

Let ${}_R M$ be a module and $\{M_i\}_{i \in I}$ a family of submodules of M . We define the *sum* of the M_i as the module

$$\sum_I M_i = \left\{ \sum_{i \in I} x_i \mid x_i \in M_i \text{ and } x_i = 0 \text{ for almost all } i \in I \right\}.$$

Clearly $\sum_I M_i \leq M$ and it is the smallest submodule of M containing all the M_i . (Notice that in the definition of $\sum_I M_i$ we need almost all the components to be zero in order to define properly the sum of elements of M).

Remark 5.1. Let ${}_R M$ be a module and $\{M_i\}_{i \in I}$ a family of submodules of M . Following the previous definitions we can construct both the module $\bigoplus_I M_i$ and module $\sum_I M_i$ (which is a submodule of M). We can define a homomorphism

$$\alpha : \bigoplus_I M_i \rightarrow \sum_I M_i, \quad (x_i)_{i \in I} \mapsto \sum_{i \in I} x_i.$$

Then $\text{Im } \alpha = \sum_I M_i$. If α is a monomorphism, then $\bigoplus_I M_i \cong \sum_I M_i$ and we say that the module $\sum_I M_i$ is the (*internal*) *direct sums* of its submodules M_i . Often we omit the word "internal" and if $M = \sum_I M_i$ and α is an isomorphism, we say that M is the direct sums of the submodules M_i and we write $M = \bigoplus_I M_i$.

Exercise 5.2. Let ${}_R M$ be a module and $\{M_i\}_{i \in I}$ a family of submodules of M . The following are equivalent:

- (1) α is an isomorphism
- (2) if $m \in \sum_I M_i$, then m can be written in a unique way as sum of elements of the M_i

6. SPLIT EXACT SEQUENCES

If L and N are R -modules, there is a short exact sequence, called *split*,

$$0 \rightarrow L \xrightarrow{i_L} L \oplus N \xrightarrow{\pi_N} N \rightarrow 0, \quad \text{with } i_L(l) = (l, 0) \quad \pi_N(l, n) = n, \quad \text{for any } l \in L, n \in N.$$

More generally:

Definition 6.1. A short exact sequence $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$ is said to be split if there is an isomorphism $M \cong L \oplus N$ such that the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & L & \xrightarrow{f} & M & \xrightarrow{g} & N \longrightarrow 0 \\ & & \parallel & & \cong \downarrow \alpha & & \parallel \\ 0 & \longrightarrow & L & \xrightarrow{i_L} & L \oplus N & \xrightarrow{\pi_N} & N \longrightarrow 0 \end{array}$$

commutes.

Proposition 6.2. *The following properties of an exact sequence $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$ are equivalent:*

- (1) *the sequence is split*
- (2) *there exists a homomorphism $\varphi : M \rightarrow L$ such that $\varphi f = \text{id}_L$*
- (3) *there exists a homomorphism $\psi : N \rightarrow M$ such that $g\psi = \text{id}_N$*

Proof. $1 \Rightarrow 2$. Since the sequence splits, then there exists α as in Definition 6.1. Let $\varphi = \pi_L \circ \alpha$. So for any $l \in L$ $\varphi f(l) = \pi_L \alpha f(l) = \pi_L(l, 0) = l$.

$1 \Rightarrow 3$ Similar (Verify!)

$2 \Rightarrow 1$. Define $\alpha : M \rightarrow L \oplus N$, $m \mapsto (\varphi(m), g(m))$. Since $\alpha f(l) = (\varphi(f(l)), g(f(l))) = (l, 0)$ and $\pi_N \alpha(m) = g(m)$ we get that the diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & L & \xrightarrow{f} & M & \xrightarrow{g} & N & \longrightarrow & 0 \\ & & \parallel & & \downarrow \alpha & & \parallel & & \\ 0 & \longrightarrow & L & \xrightarrow{i_L} & L \oplus N & \xrightarrow{\pi_N} & N & \longrightarrow & 0 \end{array}$$

commutes. Finally, by Proposition 3.2, we conclude that α is an isomorphism.

$2 \Rightarrow 3$ Similar (Verify!) □

Definition 6.3. *Given ${}_R L \leq_R M$, L is a direct summand of M if there exists a submodule ${}_R N \leq_R M$ such that M is the direct sum of L and N . N is called a complement of L . If M does not admit direct summands it is said to be indecomposable.*

By the results in the previous section, if L is a direct summand of M and N a complement of L , it means that any m in M can be written in a unique way as $m = l + n$, $l \in L$ and $n \in N$. We write $M = L \oplus N$ and $L \leq^{\oplus} M$.

Example 6.4. (1) consider the \mathbb{Z} -module $\mathbb{Z}/6\mathbb{Z}$. Then $\mathbb{Z}/6\mathbb{Z} = 3\mathbb{Z}/6\mathbb{Z} \oplus 2\mathbb{Z}/6\mathbb{Z}$. The regular module ${}_Z \mathbb{Z}$ is indecomposable

- (2) let K be a field and V a K -module. Then, by a well-know result of linear algebra, any $L \leq V$ is a direct summand of V .
- (3) Let $R = \begin{pmatrix} K & K \\ 0 & K \end{pmatrix}$. Then $R = P_1 \oplus P_2$, where $P_1 = \left\{ \begin{pmatrix} k & 0 \\ 0 & 0 \end{pmatrix} \mid k \in K \right\}$ and $P_2 = \left\{ \begin{pmatrix} 0 & k_1 \\ 0 & k_2 \end{pmatrix} \mid k_1, k_2 \in K \right\}$.

7. EXERCISES

Exercise 7.1. Let ${}_R L \leq_R M$. Show that L is a direct summand of M if and only if there exists ${}_R N \leq_R M$ such that $L + N = M$ and $L \cap N = 0$.

Exercise 7.2. Let $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$ be a split exact sequence and α the isomorphism as in Definition 6.1. Show that $M = \alpha^{-1}(L) \oplus \alpha^{-1}(N)$, $\alpha^{-1}(L) \cong L$, and $\alpha^{-1}(N) \cong N$.

8. FREE MODULES AND FINITELY GENERATED MODULES

Definition 8.1. *A module ${}_R M$ is said to be generated by a family $\{x_i\}_{i \in I}$ of elements of M if each $x \in M$ can be written as $x = \sum_I r_i x_i$, with $r_i \in R$ for any $i \in I$, and $r_i = 0$ for almost every $i \in I$.*

The $\{x_i\}_{i \in I}$ are called a set of generator of M and we write $M = \langle x_i, i \in I \rangle$.

If the coefficients r_i are uniquely determined by x , the $(x_i)_{i \in I}$ are called a basis of M .

The module M is said to be free if it admits a basis.

Proposition 8.2. *A module ${}_R M$ is free if and only if $M \cong R^{(I)}$ for some set I .*

Proof. The module $R^{(I)}$ is free with basis $(e_i)_{i \in I}$, where e_i is the canonical vector with all zero components except for the i -th equal to 1.

Conversely if M is free with basis $(x_i)_{i \in I}$, then we can define a homomorphism $\alpha : R^{(I)} \rightarrow M$, $(r_i)_{i \in I} \mapsto \sum_I r_i x_i$. It is easy to show that α is an isomorphism, as a consequence of the definition of a basis: indeed, it is clearly an epi and if $\alpha(r_i) = \sum r_i x_i = 0$, since the r_i are uniquely determined by 0, we conclude that $r_i = 0$ for all i , i.e. α is a mono. □

Given a free module M with basis $(x_i)_I$, then every homomorphism $f : M \rightarrow N$ is uniquely determined by its value on the x_i and the elements $f(x_i)$ can be chosen arbitrarily in N . Indeed, chosen the $f(x_i)$, given $x = \sum r_i x_i \in M$, we construct $f(x) = \sum r_i f(x_i)$. Since $(x_i)_{i \in I}$ is a basis this is a good definition. (Notice: analogy with vector spaces!).

Proposition 8.3. *Any module is quotient of a free module*

Proof. Let M be an R -module. Since we can always choose $I = M$, the module M admits a set of generators. Let $(x_i)_{i \in I}$ a set of generators for M and define a homomorphism $\alpha : R^{(I)} \rightarrow M$, $(r_i)_{i \in I} \mapsto \sum_i r_i x_i$. Clearly α is an epi and so $M \cong R^{(I)} / \text{Ker } \alpha$ \square

Definition 8.4. *A module ${}_R M$ is finitely generated if there exists a finite set of generators for M . A module is cyclic if it can be generated by a single element.*

By Proposition 8.3 ${}_R M$ is finitely generated if and only if there exists an epimorphism $R^n \rightarrow M$ for some $n \in \mathbb{N}$. Similarly, ${}_R M$ is cyclic if and only if $M \cong J$, for a left ideal $J \leq R$.

Example 8.5. The regular module ${}_R R$ is cyclic, generated by the unity element ${}_R R = \langle 1 \rangle$

Proposition 8.6. *Let ${}_R L \leq {}_R M$.*

- (1) *If M is finitely generated, then M/L is finitely generated.*
- (2) *If L and M/L are finitely generated, so is M*

Proof. (1) If $\{x_1, \dots, x_n\}$ is a set of generator of M , then $\{\bar{x}_1, \dots, \bar{x}_n\}$ is a set of generator for M/L .
 (2) Let $\langle x_1, \dots, x_n \rangle = L$ and $\langle \bar{y}_1, \dots, \bar{y}_m \rangle = M/L$, where $x_1, \dots, x_n, y_1, \dots, y_m \in M$. Let $x \in M$ and consider $\bar{x} = \sum_{i=1, \dots, m} r_i \bar{y}_i$ in M/L . Then $x - \sum_{i=1, \dots, m} r_i y_i \in L$ and so $x - \sum_{i=1, \dots, m} r_i y_i = \sum_{j=1, \dots, n} r_j x_j$. Hence $x = \sum_{i=1, \dots, m} r_i y_i + \sum_{j=1, \dots, n} r_j x_j$, i.e. $\{x_1, \dots, x_n, y_1, \dots, y_m\}$ is a finite set of generators of M . \square

Notice that M finitely generated doesn't imply L finitely generated. For example, let R be the ring $R = K[x_i, i \in \mathbb{N}]$. Consider the regular module ${}_R R$ and its submodule $L = \langle x_i, i \in \mathbb{N} \rangle$.

9. EXERCISES

Exercise 9.1. Show that any submodule of ${}_Z \mathbb{Z}$ is finitely generated.

Exercise 9.2. Show that the \mathbb{Z} -module \mathbb{Q} is not finitely generated.

Exercise 9.3. A module M is *simple* if $L \leq M$ implies $L = 0$ or $L = M$ (i.e. M doesn't have non trivial submodules).

- (1) show that any simple module is cyclic
- (2) Exhibit a cyclic module which is not simple.

Exercise 9.4. Let R be a ring. An element $e \in R$ is *idempotent* if $e^2 = e$. Show that

- (1) if e is idempotent, then $(1 - e)$ is idempotent and $R = Re \oplus R(1 - e)$ (where Re and $R(1 - e)$ denote the cyclic modules generated by e and $(1 - e)$, respectively)
- (2) if $R = I \oplus J$, with I and J left ideals of R , then there exist idempotents e and f such that $1 = e + f$, $I = Re$ and $J = Rf$.

10. CATEGORIES AND FUNCTORS

This is very short introduction to the basic concepts of category theory. For more details and for the set-theoretical foundation (in particular the distinction between sets and classes) we refer to S. MacLane, *Category for the working mathematician*, Graduate Texts in Math., Vol 5, Springer 1971.

Definition 10.1. A category \mathcal{C} consists in:

- (1) A class $Obj(\mathcal{C})$, called the objects of \mathcal{C} ;
- (2) for each ordered pair (C, C') of objects of \mathcal{C} , a set $Hom_{\mathcal{C}}(C, C')$ whose elements are called morphisms from C to C' ;
- (3) for each ordered triple (C, C', C'') of objects of \mathcal{C} , a map

$$Hom_{\mathcal{C}}(C, C') \times Hom_{\mathcal{C}}(C', C'') \rightarrow Hom_{\mathcal{C}}(C, C'')$$

called composition of morphisms

such that the following axioms C1, C2, C3 hold:

(before stating the axioms, we introduce the notations $\alpha : C \rightarrow C'$ for any $\alpha \in Hom_{\mathcal{C}}(C, C')$, and $\beta\alpha$ for the composition of $\alpha \in Hom_{\mathcal{C}}(C, C')$ and $\beta \in Hom_{\mathcal{C}}(C', C'')$)

- C1: if $(C, C') \neq (D, D')$, then $Hom_{\mathcal{C}}(C, C') \cap Hom_{\mathcal{C}}(D, D') = \emptyset$
- C2: if $\alpha : C \rightarrow C'$, $\beta : C' \rightarrow C''$, $\gamma : C'' \rightarrow C'''$ are morphisms, then $\gamma(\beta\alpha) = (\gamma\beta)\alpha$
- C3: for each object C there exists $1_C \in Hom_{\mathcal{C}}(C, C)$, called *identity morphism*, such that $1_C\alpha = \alpha$ and $\beta 1_C = \beta$ for any $\alpha : C' \rightarrow C$ and $\beta : C \rightarrow C'$.

Notice that, for any $C \in Obj(\mathcal{C})$, the identity morphism 1_C is unique. Indeed, if also $1'_C$ satisfies [C3], then $1_C = 1_C 1'_C = 1'_C$.

A morphism $\alpha : C \rightarrow C'$ is an *isomorphism* if there exists $\beta : C' \rightarrow C$ such that $\beta\alpha = 1_C$ and $\alpha\beta = 1_{C'}$. If α is an isomorphism, C and C' are called *isomorphic* and we write $C \cong C'$.

- Example 10.2.*
- (1) The category **Sets**: the class of objects is the class of all sets; the morphisms are the maps between sets with the usual compositions.
 - (2) The category **Ab**: the objects are the abelian groups; the morphisms are the group homomorphisms with the usual compositions.
 - (3) The category **R-Mod** for a ring R : the objects are the left R -modules and the morphisms are the module homomorphisms with the usual compositions.
 - (4) The category **Mod-R** for a ring R : the objects are the right R -modules and the morphisms are the module homomorphisms with the usual compositions.

Notice that, given a category \mathcal{C} , we can construct the *dual* category \mathcal{C}^{op} , with $Obj(\mathcal{C}^{op}) = Obj(\mathcal{C})$, $Hom_{\mathcal{C}^{op}}(C, C') = Hom_{\mathcal{C}}(C', C)$, and $\alpha*\beta = \beta\cdot\alpha$, where $*$ denotes the composition in \mathcal{C}^{op} and \cdot the composition in \mathcal{C} (\mathcal{C}^{op} is obtained from \mathcal{C} by "reversing the arrows"). Any statement regarding a category \mathcal{C} dualizes to a corresponding statement for \mathcal{C}^{op} .

Definition 10.3. Let \mathcal{B} and \mathcal{C} be two categories. A functor $F : \mathcal{B} \rightarrow \mathcal{C}$ assigns to each object $B \in \mathcal{B}$ an object $F(B) \in \mathcal{C}$, and assigns to any morphism $\beta : B \rightarrow B'$ in \mathcal{B} a morphism $F(\beta) : F(B) \rightarrow F(B')$ in \mathcal{C} , in such a way:

- F1: $F(\beta\alpha) = F(\beta)F(\alpha)$ for any $\alpha : B \rightarrow B'$, $\beta : B' \rightarrow B''$ in \mathcal{B}
- F2: $F(1_B) = 1_{F(B)}$ for any B in \mathcal{B} .

By construction, a functor $F : \mathcal{B} \rightarrow \mathcal{C}$ defines a map for any B, B' in \mathcal{B}

$$Hom_{\mathcal{B}}(B, B') \rightarrow Hom_{\mathcal{C}}(F(B), F(B')), \quad \beta \mapsto F(\beta)$$

The functor F is called *faithful* if all these maps are injective and is called *full* if they are surjective.

A functor $F : \mathcal{B}^{op} \rightarrow \mathcal{C}$ is called a *contravariant* functor from \mathcal{B} to \mathcal{C} . In particular a contravariant functor F assigns to any morphism $\beta : B \rightarrow B'$ in \mathcal{B} a morphism $F(\beta) : F(B') \rightarrow F(B)$ in \mathcal{C} .

- Example 10.4.*
- (1) Let \mathcal{B} and \mathcal{C} two categories. \mathcal{B} is a *subcategory* of \mathcal{C} if $Obj(\mathcal{B}) \subseteq Obj(\mathcal{C})$, $Hom_{\mathcal{B}}(B, B') \subseteq Hom_{\mathcal{C}}(B, B')$ for any B, B' objects of \mathcal{B} , and the compositions in \mathcal{B} and \mathcal{C} are the same. In this case there is a canonical functor $\mathcal{B} \rightarrow \mathcal{C}$ which is clearly faithful. If this functor is also full, \mathcal{B} is said a *full subcategory* of \mathcal{C} .

- (2) Let $M \in R\text{-Mod}$. As we have already observed $\text{Hom}_R(M, N)$ is an abelian group for any $N \in R\text{-Mod}$. So we can define a functor (Verify the axioms!)

$$\text{Hom}_R(M, -) : R\text{-Mod} \rightarrow \mathbf{Ab}, \quad N \mapsto \text{Hom}_R(M, N)$$

such that for any $\alpha : N \rightarrow N'$,

$$\text{Hom}_R(M, \alpha) : \text{Hom}_R(M, N) \rightarrow \text{Hom}_R(M, N'), \quad \varphi \mapsto \alpha\varphi$$

- (3) Let $M \in R\text{-Mod}$ and consider the abelian group $\text{Hom}_R(N, M)$ for any $N \in R\text{-Mod}$. So we can define a contravariant functor (Verify the axioms!)

$$\text{Hom}_R(-, M) : (R\text{-Mod})^{op} \rightarrow \mathbf{Ab}, \quad N \mapsto \text{Hom}_R(N, M)$$

such that for any $\alpha : N \rightarrow N'$,

$$\text{Hom}_R(\alpha, M) : \text{Hom}_R(N', M) \rightarrow \text{Hom}_R(N, M), \quad \psi \mapsto \psi\alpha$$

In these lectures we will deal mainly with categories having some kind of additive structure. For instance in the category $R\text{-Mod}$, any set of morphisms $\text{Hom}_R(M, N)$ is an abelian group and the composition preserves the sums.

Definition 10.5. A category \mathcal{C} is called preadditive if each set $\text{Hom}_{\mathcal{C}}(C, C')$ is an abelian group and the compositions maps $\text{Hom}_{\mathcal{C}}(C, C') \times \text{Hom}_{\mathcal{C}}(C', C'') \rightarrow \text{Hom}_{\mathcal{C}}(C, C'')$ are bilinear.

If \mathcal{B} and \mathcal{C} are preadditive categories, a functor $F : \mathcal{B} \rightarrow \mathcal{C}$ is additive if $F(\alpha + \alpha') = F(\alpha) + F(\alpha')$ for $\alpha, \alpha' : C \rightarrow C'$.

Example 10.6. The category $R\text{-Mod}$ is a preadditive category. If $M \in R\text{-Mod}$, then $\text{Hom}_R(M, -)$ and $\text{Hom}_R(-, M)$ are additive functors.

Definition 10.7. Let R and S two rings and let $F : R\text{-Mod} \rightarrow S\text{-Mod}$ be an additive functor. F is called left exact if, for any exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ in $R\text{-Mod}$, the sequence $0 \rightarrow F(L) \rightarrow F(M) \rightarrow F(N)$ in $S\text{-Mod}$ is exact. F is called right exact if, for any exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ in $R\text{-Mod}$, the sequence $F(L) \rightarrow F(M) \rightarrow F(N) \rightarrow 0$ in $S\text{-Mod}$ is exact. The functor F is exact if it is both left and right exact.

In particular, if F is exact then for any exact sequence in $R\text{-Mod}$ $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$, the corresponding sequence $0 \rightarrow F(L) \rightarrow F(M) \rightarrow F(N) \rightarrow 0$ in $S\text{-Mod}$ is exact.

Proposition 10.8. Let $X \in R\text{-Mod}$. The functor $\text{Hom}_R(X, -)$ is left exact

Proof. Let $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$ be an exact sequence in $R\text{-Mod}$. Denoted by $f^* = \text{Hom}_R(X, f)$ and $g^* = \text{Hom}_R(X, g)$, we have to show that the sequence of abelian groups $0 \rightarrow \text{Hom}_R(X, L) \xrightarrow{f^*} \text{Hom}_R(X, M) \xrightarrow{g^*} \text{Hom}_R(X, N)$ is exact. In particular, we have to show that f^* is a mono and that $\text{Im } f^* = \text{Ker } g^*$.

Let us start considering $\alpha : X \rightarrow L$ such that $f^*(\alpha) = 0$. So for any $x \in X$ $f^*(\alpha)(x) = f\alpha(x) = 0$. Since f is a mono we conclude $\alpha(x) = 0$ for any $x \in X$, that is $\alpha = 0$.

Consider now $\beta \in \text{Im } f^*$; then there exists $\alpha \in \text{Hom}_R(X, L)$ such that $\beta = f^*(\alpha) = f\alpha$. Hence $g^*(\beta) = g\beta = gf\alpha = 0$, since $gf = 0$. So we get $\text{Im } f^* \leq \text{Ker } g^*$.

Finally, let $\beta \in \text{Ker } g^*$, so that $g\beta = 0$. This means $\text{Im } \beta \leq \text{Ker } g = \text{Im } f$. For any $x \in X$ define α as $\alpha(x) = f^{-1}(\beta(x))$: α is well-defined since f is a mono and clearly $\beta = f\alpha = f^*(\alpha)$. So we get $\text{Ker } g^* \leq \text{Im } f^*$ \square

In a similar way one prove that the functor $\text{Hom}_R(-, X)$ is left exact. Notice that, since $\text{Hom}_R(-, X)$ is a contravariant functor, left exact means that for any exact sequence in $R\text{-Mod}$ $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$, the corresponding sequence of abelian groups $0 \rightarrow \text{Hom}_R(N, X) \rightarrow \text{Hom}_R(M, X) \rightarrow \text{Hom}_R(L, X)$ is exact.

Remark 10.9. Notice that if F is an additive functor and $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$ is a split exact sequence in $R\text{-Mod}$, then $0 \rightarrow F(L) \xrightarrow{F(f)} F(M) \xrightarrow{F(g)} F(N) \rightarrow 0$ is split exact. Indeed, since there exists φ such that $\varphi f = id_L$ (see Proposition 6.2), $F(\varphi)F(f) = id_{F(L)}$, so $F(f)$ is a split mono. Similarly one show that $F(g)$ is a split epi.

In particular, for a given module $X \in R\text{-Mod}$ the functors $\text{Hom}_R(X, -)$ and $\text{Hom}_R(-, X)$ could be not exact. Nevertheless, if $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is a split exact sequence in $R\text{-Mod}$, then the sequence $0 \rightarrow \text{Hom}_R(X, L) \rightarrow \text{Hom}_R(X, M) \rightarrow \text{Hom}_R(X, N) \rightarrow 0$ and the sequence $0 \rightarrow$

$\text{Hom}_R(N, X) \rightarrow \text{Hom}_R(M, X) \rightarrow \text{Hom}_R(L, X) \rightarrow 0$ are split exact. In particular $\text{Hom}_R(X, L \oplus N) = \text{Hom}_R(X, L) \oplus \text{Hom}_R(X, N)$ and $\text{Hom}_R(L \oplus N, X) = \text{Hom}_R(L, X) \oplus \text{Hom}_R(N, X)$

One often wishes to compare two functors with each other. So we introduce the following notion:

Definition 10.10. Let F and G two functors $\mathcal{B} \rightarrow \mathcal{C}$. A natural transformation $\eta : F \rightarrow G$ is a family of morphisms $\eta_B : F(B) \rightarrow G(B)$, for any $B \in \mathcal{B}$, such that for any morphism $\alpha : B \rightarrow B'$ in \mathcal{B} the following diagram in \mathcal{C} is commutative

$$\begin{array}{ccc} F(B) & \xrightarrow{\eta_B} & G(B) \\ F(\alpha) \downarrow & & \downarrow G(\alpha) \\ F(B') & \xrightarrow{\eta_{B'}} & G(B') \end{array}$$

If η_B is an isomorphism in \mathcal{C} for any $B \in \mathcal{B}$, then η is called a natural equivalence.

Two categories \mathcal{B} and \mathcal{C} are *isomorphic* if there exist functors $F : \mathcal{B} \rightarrow \mathcal{C}$ and $G : \mathcal{C} \rightarrow \mathcal{B}$ such that $GF = 1_{\mathcal{B}}$ and $FG = 1_{\mathcal{C}}$. This is a very strong notion, in fact there are several and relevant examples of categories \mathcal{B} and \mathcal{C} which have essentially the same structure, but where there is a bijective correspondence between the isomorphism classes of objects rather than between the individual objects. Therefore we define the following concept:

Definition 10.11. A functor $F : \mathcal{B} \rightarrow \mathcal{C}$ is an equivalence if there exists a functor $G : \mathcal{C} \rightarrow \mathcal{B}$ and natural equivalences $GF \rightarrow 1_{\mathcal{B}}$ and $FG \rightarrow 1_{\mathcal{C}}$

If the functor F is contravariant and gives an equivalence between \mathcal{B}^{op} and \mathcal{C} , we say that F is a *duality*.

Proposition 10.12. A functor $F : \mathcal{B} \rightarrow \mathcal{C}$ is an equivalence if and only if it is full and faithful, and every object of \mathcal{C} is isomorphic to an object of the form $F(B)$, with $B \in \mathcal{B}$.

11. EXERCISE

Exercise 11.1. Let (P, \leq) be a partially ordered set. Let us define a category \mathcal{C} in this way: the objects of \mathcal{C} are the elements of P , and with a unique morphism $p \rightarrow q$ whenever $p \leq q$, while $\text{Hom}_{\mathcal{C}}(p, q) = 0$ if $p \not\leq q$. Verify that the axioms [C1], [C2], [C3] are satisfied. This is an example of a *small* category, i.e. a category where the class of objects is a set.

Exercise 11.2. Let $\varphi : R \rightarrow S$ be a homomorphism of rings. Each left S -module M has also a structure of left R -module, defining $rx := \varphi(r)x$ for any $x \in M$ and any $r \in R$. Let $\varphi^* : S\text{-Mod} \rightarrow R\text{-Mod}$, $M \mapsto M$, $\alpha \mapsto \alpha$ for any $M \in S\text{-Mod}$ and for any $\alpha \in \text{Hom}_S(M, N)$. Verify that φ^* is an additive and faithful functor (called *restriction of scalars*)

Exercise 11.3. A functor F is exact if and only if $F(L) \rightarrow F(M) \rightarrow F(N)$ is exact whenever $L \rightarrow M \rightarrow N$ is exact.

12. PROJECTIVE MODULES

In general, for a given R -module M , the functor $\text{Hom}_R(M, -)$ is left exact but not right exact. In this section we study the R -modules P for which $\text{Hom}_R(P, -)$ is also right exact.

Definition 12.1. A module $P \in R\text{-Mod}$ is projective if $\text{Hom}_R(P, -)$ is an exact functor.

The right exactness is equivalent to require that for any $M \xrightarrow{g} N \rightarrow 0$ in $R\text{-Mod}$ the homomorphism $\text{Hom}_R(P, M) \xrightarrow{\text{Hom}_R(P, g)} \text{Hom}_R(P, N)$ is an epi, that is for any $\varphi \in \text{Hom}_R(P, N)$ there exists $\psi \in \text{Hom}_R(P, M)$ such that $g\psi = \varphi$.

$$\begin{array}{ccc} M & \xrightarrow{g} & N & \longrightarrow & 0 \\ & \swarrow \psi & \uparrow \varphi & & \\ & & P & & \end{array}$$

Example 12.2. Any free module is projective. Indeed, let $R^{(I)}$ a free R -module with $(x_i)_{i \in I}$ a basis. Given $M \xrightarrow{g} N \rightarrow 0$ and $\varphi : R^{(I)} \rightarrow N$ in $R\text{-Mod}$, let $m_i \in M$ such that $g(m_i) = \varphi(x_i)$ for any $i \in I$. Define $\psi(x_i) = m_i$ and, for $x = \sum r_i x_i$, $\psi(x) = \sum r_i m_i$. We get that $g\psi = \varphi$. Notice that from the construction is clear that the homomorphism ψ could be not unique.

Proposition 12.3. Let $P \in R\text{-Mod}$. The following are equivalent:

- (1) P is projective
- (2) P is a direct summand of a free module
- (3) every exact sequence $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} P \rightarrow 0$ splits.

Proof. $1 \Rightarrow 3$ Let $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} P \rightarrow 0$ be an exact sequence in $R\text{-Mod}$ and consider the homomorphism $1_P : P \rightarrow P$. Since P is projective there exists $\psi : P \rightarrow M$ such that $g\psi = 1_P$. By Proposition 6.2 we conclude that the sequence splits.

$3 \Rightarrow 2$ The module P is a quotient of a free module, so there exist an exact sequence $0 \rightarrow K \xrightarrow{f} R^{(I)} \xrightarrow{g} P \rightarrow 0$, which is split.

$2 \Rightarrow 1$ If $R^{(I)} = P \oplus L$, then $\text{Hom}_R(R^{(I)}, N) = \text{Hom}_R(P, N) \oplus \text{Hom}_R(L, N)$ for any $N \in R\text{-Mod}$. So let us consider the homomorphisms

$$\begin{array}{ccc} M & \xrightarrow{g} & N & \longrightarrow & 0 \\ & & \uparrow \varphi & & \\ & & P & & \end{array} \quad \text{and} \quad \begin{array}{ccc} M & \xrightarrow{g} & N & \longrightarrow & 0 \\ & \swarrow \alpha & \uparrow (\varphi, 0) & & \\ & & R^{(I)} & & \end{array}$$

where $(\varphi, 0)(p + l) = \varphi(p) + 0(l) = \varphi(p)$ for any $p \in P$ and $l \in L$ and α exists since $R^{(I)}$ is projective. Then $\alpha = (\psi, \beta)$, with $\psi \in \text{Hom}_R(P, N)$ and $\beta \in \text{Hom}_R(L, N)$, where $\alpha(p + l) = \psi(p) + \beta(l)$ for any $p \in P$ and $l \in L$. Hence $g(\psi(p)) = g(\alpha(p)) = \varphi(p)$ for any $p \in P$. So we conclude that P is projective. \square

Example 12.4. (1) Let R be a principal ideal domain (for instance, $R = \mathbb{Z}$). Then any projective module is free. In particular, free abelian groups and projective abelian group coincide.

- (2) Let $R = \mathbb{Z}/6\mathbb{Z}$. Then $\mathbb{Z}/6\mathbb{Z} = 3\mathbb{Z}/6\mathbb{Z} \oplus 2\mathbb{Z}/6\mathbb{Z}$. The ideals $3\mathbb{Z}/6\mathbb{Z}$ and $2\mathbb{Z}/6\mathbb{Z}$ are projective R -modules, but not free R -modules (why?)

Proposition 12.5. Let $P \in R\text{-Mod}$. P is projective if and only if there exists a family $(\varphi_i, x_i)_{i \in I}$ with $\varphi_i \in \text{Hom}_R(P, R)$ and $x_i \in P$ such that for any $x \in P$ one has $x = \sum_i \varphi_i(x)x_i$ where $\varphi_i(x) = 0$ for almost every $i \in I$.

Proof. Let P be projective and let $R^{(I)} \xrightarrow{\beta} P \rightarrow 0$ be a split epi. Consider $(e_i)_{i \in I}$ a basis of $R^{(I)}$ and define $x_i = \beta(e_i)$. Observe that $\beta(\sum_i r_i e_i) = \sum_i r_i \beta(e_i) = \sum_i r_i x_i$. By Proposition 6.2, there exists $\varphi : P \rightarrow R^{(I)}$ such that $\beta\varphi = id_P$, which induces homomorphisms $\varphi_i = \pi_i \varphi$ where π_i is the projection on the i -th component, so $\varphi_i(x) \in R$ for any $i \in I$ and $\varphi(x) = \sum \varphi_i(x)$. Hence for any $x \in P$ one has $x = \beta\varphi(x) = \beta(\sum_i \varphi_i(x)) = \sum_i \varphi_i(x)x_i$, so $(\varphi_i, x_i)_{i \in I}$ satisfies the stated properties.

Conversely, let $(\varphi_i, x_i)_{i \in I}$ satisfy the statement and let $\beta : R^{(I)} \rightarrow P$, $e_i \mapsto x_i$. The homomorphism β is an epi, since the family $(x_i)_{i \in I}$ generates P , and $\beta(\sum r_i) = \sum r_i x_i$. Define $\varphi : P \rightarrow R^{(I)}$, $x \mapsto \sum \varphi_i(x)$. Then for any $x \in P$ one gets $\beta\varphi(x) = \beta(\sum \varphi_i(x)) = \sum \varphi_i(x)x_i = x$. By Proposition 6.2 we conclude that β is a split epi and so P is projective. \square

Note that, from the results in the previous sections, the projective module ${}_R R$ plays a crucial role in the category $R\text{-Mod}$, since for any $M \in R\text{-Mod}$ there exists an epi $R^{(I)} \rightarrow M \rightarrow 0$, for some set I . A module with such property is called a *generator* and so R is a *projective generator* for $R\text{-Mod}$.

In particular, for any $M \in R\text{-Mod}$ there exists a short exact sequence $0 \rightarrow K \rightarrow P_0 \rightarrow M \rightarrow 0$, with P_0 projective. The same holds for the module K , and so, iterating the argument, we can construct an exact sequence

$$\cdots \rightarrow P_i \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$$

where all the P_i are projective. Such a sequence is called a *projective resolution* of P . It is clearly not unique.

It is natural to ask if, for a given $M \in R\text{-Mod}$, there exists a projective module P and a "minimal" epi $P \rightarrow M \rightarrow 0$, in the sense that $f|_L : L \rightarrow M$ is epi for no proper projective submodule of P . More precisely, we define:

Definition 12.6. A homomorphism $f : M \rightarrow N$ is right minimal if for any $g \in \text{End}_R(M)$ such that $fg = f$, one gets g is an isomorphism.

If P_M is a projective module and $P_M \rightarrow M$ is epimorphism right minimal, then P_M is a projective cover of M .

Remark 12.7. Consider the diagram

$$\begin{array}{ccc} & & 0 \\ & & \uparrow \\ P_M & \xrightarrow{f} & M \longrightarrow 0 \\ & \swarrow \psi & \uparrow g \\ & & P \end{array}$$

where P_M is a projective cover of M and P is a projective module. Since P_M and P are projective, there exist φ and ψ such that the diagram commutes. Hence $f\psi = g$ and $g\varphi = f$, so $f\psi\varphi = f$ and, since f is a right minimal, we conclude $\psi\varphi$ is an iso. In particular φ is a mono. Define $\theta : P \rightarrow P_M$ as $\theta = (\psi\varphi)^{-1}\psi$: then $\theta\varphi = id_P$ and so φ is a split mono (see Proposition 6.2). We conclude that P_M is a direct summand of P . This explains the minimality property of the projective cover announced above.

If also P is a projective cover of M , using the same argument we get that $\varphi\psi$ is an iso, that is $\varphi = \psi^{-1}$ and P_M is isomorphic to P . We have shown that the projective cover is unique (modulo isomorphisms).

We state the following characterization of projective covers:

Theorem 12.8. Let P a projective module. Then $P \xrightarrow{f} M \rightarrow 0$ is a projective cover of M if and only if $\text{Ker } f$ is a superfluous submodule of P (i.e. for any submodule $L \leq P$, $L + \text{Ker } f = P$ implies $L = P$.)

Observe that, given $M \in R\text{-Mod}$, a projective cover for M could not exist. A ring in which any module admits a projective cover is called *semiperfect*

Let now $M \in R\text{-Mod}$ and suppose there exist a projective resolution of M

$$\cdots P_2 \xrightarrow{f_2} P_1 \xrightarrow{f_1} P_0 \xrightarrow{f_0} M \rightarrow 0$$

such that P_0 is a projective cover of M and P_i is a projective cover of $\text{Ker } f_{i-1}$ for any $i \in \mathbb{N}$. Such a resolution is called a *minimal projective resolution* of M .

13. EXERCISE

Exercise 13.1. Let $P_1, P_2, \dots, P_n \in R\text{-Mod}$. Then $\bigoplus_{i=1, \dots, n} P_i$ is projective if and only if P_i is projective for any $i = 1, \dots, n$.

Exercise 13.2. Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ a short exact sequence in $R\text{-Mod}$. If L and N are projective, then M is projective

Exercise 13.3. Show that any abelian group $n\mathbb{Z}$, $n \in \mathbb{N}$, is a projective \mathbb{Z} -module.

14. BIMODULES

Definition 14.1. Let R and S rings. An abelian group M is a left R -right S -bimodule if M is a left R -module and a right S -module such that the two scalar multiplications satisfy $r(xs) = (rx)s$ for any $r \in R$, $s \in S$, $x \in M$. We write ${}_R M_S$.

Example 14.2. Let $M \in R\text{-Mod}$ and consider $S = \text{End}_R^r(M)$, the ring of homomorphism R -linear of M , where homomorphisms act on the right (i.e. $mf = f(m)$ and $m(fg) = g(f(m))$). So M is a right S -module (Verify!) and ${}_R M_S$ is a bimodule. Indeed $(rm)f = f(rm) = rf(m) = r(f(m))$ for any $r \in R$, $m \in M$ and $f \in S$.

Given a bimodule ${}_R M_S$ and a left R -module N , the abelian group $\text{Hom}_R(M, N)$ is naturally endowed with a structure of left S -module, by defining $(sf)(x) := f(xs)$ for any $f \in \text{Hom}_R(M, N)$ and any $x \in M$. (Verify! crucial point: $(s_1(s_2f))(x) = (s_2f(xs_1)) = f(xs_1s_2) = ((s_1s_2)f)(x)$).

Similarly, if ${}_R N_T$ is a left R -right T -bimodule and $M \in R\text{-Mod}$, then $\text{Hom}_R(M, N)$ is naturally endowed with a structure of right T -module, by defining $(ft)(x) := f(x)t$ (Verify! crucial point: $(f(t_1t_2))(x) = f(x)(t_1t_2) = (f(x))t_1t_2 = ((ft_1)(x))t_2 = ((ft_1)t_2)(x)$).

Moreover, one can show that if ${}_R M_S$ and ${}_R N_T$ are bimodules, then $\text{Hom}_R({}_R M_S, {}_R N_T)$ is a left S -right T -bimodule (Verify!).

Arguing in a similar way for right R -modules, if ${}_S M_R$ and ${}_T N_R$ are bimodules, then the abelian group $\text{Hom}_R({}_S M_R, {}_T N_R)$ is a left T -right S -bimodule, by $(tf)(x) = t(f(x))$ and $(fs)(x) = f(sx)$.

15. INJECTIVE MODULES

In this section we study the R -modules E for which $\text{Hom}_R(-, E)$ is an exact functor. Observe that many results we are going to show are dual of those proved for projective modules.

Definition 15.1. A module $E \in R\text{-Mod}$ is injective if $\text{Hom}_R(-, E)$ is an exact functor.

The exactness is equivalent to require that for any $0 \rightarrow L \xrightarrow{f} M$ in $R\text{-Mod}$ the homomorphism $\text{Hom}_R(M, E) \xrightarrow{\text{Hom}_R(f, E)} \text{Hom}_R(L, E)$ is an epi, that is for any $\varphi \in \text{Hom}_R(L, E)$ there exists $\psi \in \text{Hom}_R(M, E)$ such that $\psi f = \varphi$.

$$\begin{array}{ccccc} 0 & \longrightarrow & L & \xrightarrow{f} & M \\ & & \downarrow \varphi & \swarrow \psi & \\ & & E & & \end{array}$$

Any module is quotient of a projective module. Does the dual property hold? that is, given any module $M \in R\text{-Mod}$, is it true that M embeds in a injective R -module? In the sequel we will answer to this crucial question.

An abelian group G is *divisible* if, for any $n \in \mathbb{Z}$ and for any $g \in G$, there exists $t \in G$ such that $g = nt$. We are going to show that an abelian group is injective if and only if it is divisible. We need the the following useful criterion to check whether a module is injective, known as Baer's Lemma.

Lemma 15.2. Let $E \in R\text{-Mod}$. The module E is injective if and only if for any left ideal J of R and for any $\varphi \in \text{Hom}_R(J, E)$ there exists $\psi \in \text{Hom}_R(R, E)$ such that $\psi i = \varphi$, where i is the canonical inclusion $0 \rightarrow J \xrightarrow{i} R$.

The lemma states that it is sufficient to check the injectivity property only for left ideals of the ring. In particular, the Baer's Lemma says that E is injective if and only if for any $RJ \leq {}_R R$ and for any $\varphi \in \text{Hom}_R(J, E)$ there exists $y \in E$ such that $\varphi(x) = xy$ for any $x \in J$.

Proposition 15.3. A module $G \in \mathbb{Z}\text{-Mod}$ is injective if and only if it is divisible.

Proof. Let us assume G injective, consider $n \in \mathbb{Z}$ and $g \in G$ and the commutative diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & \mathbb{Z}n & \xrightarrow{i} & \mathbb{Z} \\ & & \downarrow \varphi & \swarrow \psi & \\ & & G & & \end{array}$$

where $\varphi(sn) = sg$ for any $s \in \mathbb{Z}$ and ψ exists since G is injective. Let $t = \psi(1)$, $t \in G$. Then $\varphi(n) = \psi(i(n))$ implies $g = nt$ and we conclude that G is divisible.

Conversely, suppose G divisible and apply Baer's Lemma. The ideal of \mathbb{Z} are of the form $\mathbb{Z}n$ for $n \in \mathbb{Z}$, so we have to verify that for any $\varphi \in \text{Hom}_{\mathbb{Z}}(\mathbb{Z}n, G)$ there exists ψ such that

$$\begin{array}{ccccc} 0 & \longrightarrow & \mathbb{Z}n & \xrightarrow{i} & \mathbb{Z} \\ & & \downarrow \varphi & \swarrow \psi & \\ & & G & & \end{array}$$

commutes. Let $g \in G$ such that $\varphi(n) = g$. Since \mathbb{Z} is a free \mathbb{Z} -module, define $\psi(1) = t$ where $g = nt$ and so $\psi(r) = rt$ for any $r \in \mathbb{Z}$. Hence $\varphi(sn) = sg = snt = \psi(i(sn))$. \square

The result stated in the previous proposition holds for any Principal Ideal Domain R (see Exercise 16.1).

Example 15.4. The \mathbb{Z} -module \mathbb{Q} is injective.

Remark 15.5. Any abelian group G embeds in a injective abelian group. Indeed, consider a short exact sequence $0 \rightarrow K \rightarrow \mathbb{Z}^{(I)} \rightarrow G \rightarrow 0$ and the canonical inclusion in $\mathbb{Z}\text{-Mod}$ $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q}$. One easily check that $\mathbb{Q}^{(I)}/K$ is divisible (Verify!) and so injective. Then we get the induced monomorphism $0 \rightarrow G \cong \mathbb{Z}^{(I)}/K \rightarrow \mathbb{Q}^{(I)}/K$.

Proposition 15.6. *Let R be a ring. If $D \in \mathbb{Z}\text{-Mod}$ is injective, then $\text{Hom}_{\mathbb{Z}}(R, D)$ is an injective left R -module*

Proof. First notice that, since ${}_{\mathbb{Z}}R_R$ is a bimodule, $\text{Hom}_{\mathbb{Z}}(R, D)$ is naturally endowed with a structure of left R -module. In order to verify that it is injective, we apply Baer's Lemma. So let ${}_R I \leq {}_R R$ and $h : I \rightarrow \text{Hom}_{\mathbb{Z}}(R, D)$ an R -homomorphism. Then $\gamma : I \rightarrow D$, $a \mapsto h(a)(1)$ defines a \mathbb{Z} -homomorphism and, since D is an injective abelian group, there exists $\bar{\gamma} : R \rightarrow D$ which extends γ . Now we have, for any $a \in I$ and $r \in R$,

$$(a\bar{\gamma})(r) = \bar{\gamma}(ra) = \gamma(ra) = [h(ra)](1) = [rh(a)](1) = [h(a)](r)$$

so $h(a) = a\bar{\gamma}$ for any $a \in I$. Hence we conclude $\text{Hom}_{\mathbb{Z}}(R, D)$ is injective by Baer's Lemma. \square

Corollary 15.7. *Let $M \in R\text{-Mod}$. Then there exists an injective module $E \in R\text{-Mod}$ and a monomorphism $0 \rightarrow M \rightarrow E$.*

Proof. Consider the isomorphism of \mathbb{Z} -modules $\varphi : \text{Hom}_R(R, M) \rightarrow M$, $f \mapsto f(1)$. Observe that since ${}_R R_R$ is a left R -right R -bimodule, then $\text{Hom}_R(R, M)$ is naturally endowed with a structure of left R -module. One easily check that φ is also R -linear, hence ${}_R M \cong \text{Hom}_R(R, M) \leq \text{Hom}_{\mathbb{Z}}(R, M)$. By Remark 15.5, there is a mono of \mathbb{Z} -modules $0 \rightarrow M \rightarrow G$ from which we obtain a mono of R -modules $0 \rightarrow \text{Hom}_{\mathbb{Z}}(R, M) \rightarrow \text{Hom}_{\mathbb{Z}}(R, G)$, where $\text{Hom}_{\mathbb{Z}}(R, G)$ is an injective left R -module by Proposition 15.6. \square

Since any module M embeds in a injective one, it is natural to ask whether there exists a "minimal" injective module containing M .

Definition 15.8. *A homomorphism $f : M \rightarrow N$ is left minimal if for any $g \in \text{End}_R(N)$ such that $gf = f$, one gets g is an isomorphism.*

If E_M is an injective module and $M \rightarrow E_M$ is a monomorphism left minimal, then E_M is an injective envelope of M .

Remark 15.9. Consider the diagram

$$\begin{array}{ccccc} & & 0 & & \\ & & \downarrow & & \\ 0 & \longrightarrow & M & \xrightarrow{f} & E_M \\ & & \downarrow g & \searrow \psi & \nearrow \gamma \\ & & E & \xleftarrow{\varphi} & \end{array}$$

where E_M is an injective envelope of M and E is an injective module. Since E_M and E are injective, there exist φ and ψ such that the diagram commutes. Hence $\psi g = f$ and $\varphi f = g$, so $\psi\varphi f = f$ and, since f is left minimal, we conclude that $\psi\varphi$ is an iso. In particular φ is a mono and so it is a split mono. We conclude that E_M is a direct summand of E . This explains the minimality property of the injective envelope announced above.

If also E is an injective envelope of M , using the same argument we get that $\varphi\psi$ is an iso, that is ψ is an iso and E_M is isomorphic to E . We have shown that the injective envelope is unique (modulo isomorphisms).

We state the following characterization of injective envelope.

Theorem 15.10. *Let E be an injective module. Then $0 \rightarrow M \xrightarrow{f} E$ is an injective envelope if and only if $\text{Im } f$ is an essential submodule of E (i.e. for any submodule $L \leq E$, $L \cap \text{Im } f \neq \{0\}$)*

Proof. Suppose $0 \rightarrow M \xrightarrow{f} E$ is an injective envelope and let $L \leq E$ such that $L \cap \text{Im } f = \{0\}$. Then $\text{Im } f \oplus L \leq E$ and we can consider the commutative diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & M & \xrightarrow{f} & \text{Im } f \oplus L & \xrightarrow{i} & E \\ & & \downarrow f & \nearrow (\text{id}, 0) & \searrow \varphi & \nearrow \varphi & \\ & & E & & & & \end{array}$$

where i is the canonical inclusion of $\text{Im } f \oplus L$ in E and φ exists since E is injective. Then $\varphi f = f$ but φ is clearly not an iso.

Conversely, let $\text{Im } f$ be essential in M and let $g \in \text{End}_R(E)$ such that $gf = f$. Since f is an essential mono we conclude that g is a mono (see Exercise 16.4), so it is a split mono. In particular, $\text{Im } f \leq \text{Im } g \leq E$, contradicting the essentiality of $\text{Im } f$. \square

Not every module has a projective cover. Thus the next result is especially remarkable

Theorem 15.11. *Every module has an injective envelope.*

Proof. Let $M \in R\text{-Mod}$; by Corollary 15.7 there exists an injective module Q such that $0 \rightarrow M \rightarrow Q$. Consider the set $\{E' \mid M \leq E' \leq Q \text{ and } M \text{ essential in } E'\}$. One easily check that it is an inductive set so, by Zorn's Lemma, it contains a maximal element E . Let us show that E is a direct summand of Q and so E is injective (see Exercise 16.3). To this aim, consider the set $\{F' \mid F' \leq Q \text{ and } F' \cap E = 0\}$. It is inductive so, again by Zorn's Lemma, it contains a maximal element F . Then there exists an obvious iso $g : E \oplus F/F \rightarrow E$ and $E \oplus F/F \leq Q/F$: from the maximality of F it follows that $E \oplus F/F \leq Q/F$ is an essential inclusion (Verify!) so consider the diagram

$$\begin{array}{ccc} 0 & \longrightarrow & E \oplus F/F & \xrightarrow{j} & Q/F \\ & & \downarrow g & \swarrow \varphi & \\ & & Q & & \end{array}$$

where j is the canonical inclusion and φ exists since Q is injective. Moreover φ is a mono since $\varphi j = g$ is a mono and j is an essential mono (see Exercise 16.4). It follows that M is essential in $E = \text{Im } g$ and $E = \text{Im } g = \varphi(E \oplus F/F)$ is essential in $\text{Im } \varphi$. Thus M is essential in $\text{Im } \varphi$ so, from the maximality of E we conclude that $E = \text{Im } \varphi$ and hence $\varphi(E \oplus F/F) = \varphi(Q/F)$. Since φ is a mono we conclude $E \oplus F = Q$. \square

Proposition 15.12. *Let $E \in R\text{-Mod}$. The following are equivalent:*

- (1) E is injective
- (2) every exact sequence $0 \rightarrow E \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$ splits.

Proof. $1 \Rightarrow 2$ Consider the commutative diagram

$$\begin{array}{ccc} 0 & \longrightarrow & E & \xrightarrow{f} & M \\ & & \downarrow \text{id}_E & \swarrow \varphi & \\ & & E & & \end{array}$$

where φ exists since E is injective. Since $\varphi f = \text{id}_E$, by Proposition 6.2 we conclude that f is a split mono.

$2 \Rightarrow 1$ By Corollary 15.7 there exists an exact sequence $0 \rightarrow E \rightarrow F \rightarrow N \rightarrow 0$, where F is an injective module. Since the sequence splits, we get that E is a direct summand of a injective module, and so E is injective (see Exercise 16.3). \square

Comparing the previous proposition with the analogous one for projective modules (see Proposition 12.3), there is an evident difference. Speaking about projective modules, we saw that a special role is played by the projective generator R . Does a module with the dual property exist? An injective module $E \in R\text{-Mod}$ such that any $M \in R\text{-Mod}$ embeds in E^{I_M} , for a set I_M , is called an *injective cogenerator* of $R\text{-Mod}$. We will see in the sequel that such a module always exists.

Remark 15.13. Dualizing what we showed in the projective case, for any module $M \in R\text{-Mod}$ there exists a long exact sequence $0 \rightarrow M \xrightarrow{f_3} E_0 \xrightarrow{f_1} E_1 \xrightarrow{f_3} E_2 \rightarrow \dots$, where the E_i are injective. This is called an *injective coresolution* of M . If E_0 is an injective envelope of M and E_i in an injective envelope of $\text{Ker } f_i$ for any $i \geq 1$, then the sequence is called a *minimal injective coresolution* of M .

16. EXERCISES

Exercise 16.1. Let R be a Principal Ideal Domain. Prove that an R -module is injective if and only if it is divisible.

Exercise 16.2. Let G be a divisible abelian group. Then $G^{(I)}$ and G/N are divisible, for any set I and for any subgroup N of G .

Exercise 16.3. Let E_i for $i = 1, \dots, n$ in R -Mod. Then $\bigoplus_{i \in I} E_i$ is injective if and only if E_i is injective for any $i = 1 \dots n$.

Exercise 16.4. A monomorphism $0 \rightarrow L \rightarrow M$ in R -Mod is called *essential monomorphism* if $\text{Im } L$ is essential in M . Prove that if f is an essential morphism and gf is a mono, then g is a mono.

Exercise 16.5. Let $0 \rightarrow M \xrightarrow{f} L$ and $0 \rightarrow L \xrightarrow{g} N$ two essential monomorphism. Show that gf is an essential monomorphism.

17. ON THE LATTICE OF SUBMODULES OF M

Let $M \in R\text{-Mod}$ and consider the partially ordered set $\mathcal{L}_M = \{L \mid L \leq M\}$. Then \mathcal{L}_M is a complete lattice, where for any $N, L \in \mathcal{L}$, $\sup\{N, L\} = L + N$ and $\inf\{N, L\} = L \cap N$. The greatest element of \mathcal{L}_M is M and the smallest is $\{0\}$.

Given an arbitrary module $M \in R\text{-Mod}$, it is natural to ask whether minimal or maximal elements of \mathcal{L} exist. They are exactly the maximal submodules of M and the simple submodules of M , respectively. More precisely we introduce the following definitions:

Definition 17.1. A module $S \in R\text{-Mod}$ is simple if $L \leq S$ implies $L = \{0\}$ or $L = S$.

A submodule $N \leq M$ is a maximal submodule of M if $N \leq L \leq M$ implies $L = N$ or $L = M$.

Example 17.2. (1) Let K be a field. Then K is the unique (modulo isomorphisms) simple module in $K\text{-Mod}$.

(2) In $\mathbb{Z}\text{-Mod}$ any abelian group $\mathbb{Z}/\mathbb{Z}p$ with p prime is a simple abelian group. So in $\mathbb{Z}\text{-Mod}$ there are infinite simple modules.

(3) The regular module \mathbb{Z} does not contain any simple submodule, since any ideal of \mathbb{Z} is of the form $\mathbb{Z}n$ and $\mathbb{Z}m \leq \mathbb{Z}n$ whenever n divides m .

In general, it is not true that any module contains a simple or a maximal submodule. Nevertheless we have the following result (see also Exercise 18.1)

Proposition 17.3. Let R be a ring and ${}_R I \leq {}_R R$. There exists a maximal left ideal M of R such that $I \leq M \leq R$. In particular R admits maximal left ideals.

Proof. Let $\mathcal{F} = \{L \mid I \leq L < R\}$. The set \mathcal{F} is inductive since, given a sequence $L_0 \leq L_1 \leq \dots$, the left ideal $\bigcup L_i$ contains all the L_i and it is a proper ideal of R . Indeed, if $\bigcup L_i = R$, there would exist an index $j \in \mathbb{N}$ such that $1 \in L_j$ and so $L_j = R$. So by Zorn's Lemma, \mathcal{F} has a maximal element, which is clearly a maximal left ideal of R . \square

Example 17.4. Consider the regular module \mathbb{Z} . Then $\mathbb{Z}p$ is a maximal submodule of \mathbb{Z} for any prime number p . Moreover the ideal $\mathbb{Z}n$ is contained in $\mathbb{Z}p$ for any p such that $p \mid n$.

Remark 17.5. Let $\mathcal{M} \leq R$ a maximal left ideal of R . Clearly R/\mathcal{M} is a simple R -module, and this shows that simple modules always exist in $R\text{-Mod}$, for any ring R .

Conversely, let $S \in R\text{-Mod}$ be a simple module. So $S = Rx$ for an element $x \in S$ and let $\text{Ann}_R(x) = \{r \in R \mid rx = 0\}$. $\text{Ann}_R(x)$ is a maximal left ideal of R , since it is the kernel of the epimorphism $\varphi : R \rightarrow S, 1 \mapsto x$, and hence $S \cong R/\text{Ann}_R(x)$.

Finally, for any simple module S consider the module $\text{Ann}_R(S) = \bigcap_{x \in S} \text{Ann}_R(x)$. It is easy to show that $\text{Ann}_R(S)$ is a two-sided ideal of R , called the *annihilator of the simple module S* (see Exercise 18.2).

The simple modules play an crucial role in the study of the category $R\text{-Mod}$, for instance:

Proposition 17.6. Let $E \in R\text{-Mod}$ be an injective module. The module E is a cogenerator of $R\text{-Mod}$ if and only if for any simple module $S \in R\text{-Mod}$ there exists a mono $0 \rightarrow S \rightarrow E^{I_S}$, for a set I_S .

Proof. Assume for any simple module $S \in R\text{-Mod}$ there exists a mono $0 \rightarrow S \xrightarrow{f_S} E^{I_S}$, for a set I_S . Then there exist $j \in I_S$ such that $\pi_j \circ f : S \rightarrow E$ is not the zero map. So, since $\text{Ker}(\pi_j \circ f) \leq S$, we get that for any simple module S there exists a mono $\pi_j \circ f : S \rightarrow E$. Let now $M \in R\text{-Mod}$, and $x \in M, x \neq 0$. So $Rx \leq M$ and $Rx \cong R/\text{Ann}_R(x)$. By Proposition 17.3 there exists a maximal submodule $\mathcal{M} \leq R$ such that $\text{Ann}_R(x) \leq \mathcal{M}$. Consider the diagram

$$\begin{array}{ccccc}
 0 & \longrightarrow & Rx \cong R/\text{Ann}_R(x) & \longrightarrow & M \\
 & & \downarrow & & \searrow \\
 & & R/\mathcal{M} \cong S & & \varphi_x \\
 & & \downarrow & & \swarrow \\
 & & E & &
 \end{array}$$

where $\varphi_x : M \rightarrow E$ exists since E is injective. In particular $\varphi_x(x) \neq 0$. Hence we can construct a mono $\varphi : M \rightarrow E^M, x \mapsto (0, 0, \dots, 0, \varphi_x(x), 0, \dots, 0)$, where $\varphi_x(x)$ is the x^{th} position. \square

Corollary 17.7. *Let $\{S_\lambda\}_{\lambda \in \Lambda}$ be a set of representative of the simple modules (modulo isomorphisms) in $R\text{-Mod}$. Then the injective envelope $E(\oplus S_\lambda)$ is a minimal injective cogenerator of $R\text{-Mod}$*

Proof. The injective module $E(\oplus S_\lambda)$ cogenerates all the simple modules, so by the previous Proposition it is an injective cogenerator. If W is a injective cogenerator of $R\text{-Mod}$, since $S_\lambda \leq W$ for any $\lambda \in \Lambda$ (see the argument in the previous proof) one gets $\oplus S_\lambda \leq W$. Since $E(\oplus S_\lambda)$ is the injective envelope of $\oplus S_\lambda$, we conclude $E(\oplus S_\lambda) \overset{\oplus}{\leq} W$. \square

Remark 17.8. If there is a finite number of simple modules in $R\text{-Mod}$ (modulo isomorphisms), S_1, S_2, \dots, S_n , then $E(\oplus S_i) = \oplus E(S_i)$ is a minimal injective cogenerator of $R\text{-Mod}$

Definition 17.9. *Let $M \in R\text{-Mod}$. The socle of M is the submodule $\text{Soc}(M) = \sum\{S \mid S \text{ is a simple submodule of } M\}$. The radical of M is the submodule $\text{Rad}(M) = \cap\{N \mid N \text{ is a maximal submodule of } M\}$.*

Remark 17.10. If M does not contain any simple module, we set $\text{Soc}(M) = 0$. If M does not contain any maximal submodule, we set $\text{Rad}(M) = M$.

In the next Proposition we list some important properties of the socle and of the radical of a module. We leave the proofs for exercise.

Proposition 17.11. *Let $M \in R\text{-Mod}$.*

- (1) $\text{Soc}(M) = \oplus\{S \mid S \text{ is a simple submodule of } M\}$. In particular, $\text{Soc}(M)$ is a semisimple module.
- (2) $\text{Soc}(M) = \cap\{L \mid L \text{ is an essential submodule of } M\}$.
- (3) $\text{Rad}(M) = \sum\{U \mid U \text{ is a superfluous submodule of } M\}$.
- (4) Let $f : M \rightarrow N$. Let $f(\text{Soc}(M)) \leq \text{Soc}(N)$ and $f(\text{Rad}(M)) \leq \text{Rad}(N)$.
- (5) if $M = \oplus_{\lambda \in \Lambda} M_\lambda$, then $\text{Soc}(M) = \oplus_{\lambda \in \Lambda} \text{Soc}(M_\lambda)$ and $\text{Rad}(M) = \oplus_{\lambda \in \Lambda} \text{Rad}(M_\lambda)$.
- (6) $\text{Rad}(M/\text{Rad}(M)) = 0$ and $\text{Soc}(\text{Soc}(M)) = \text{Soc}(M)$.
- (7) If M is finitely generated, then $\text{Rad}(M)$ is a superfluous submodule of M .

Remark 17.12. It is clear that the radical can be described also by

$$\text{Rad}(M) = \{x \in M \mid \varphi(x) = 0 \text{ for every } \varphi : M \rightarrow S \text{ with } S \text{ simple}\}$$

Indeed, given $\varphi : M \rightarrow S$ with S simple, the kernel of φ is a maximal submodule of M . Conversely, if N is a maximal submodule of M , then consider $\pi : M \rightarrow M/N$ where M/N is simple.

A crucial role is played by the radical of the regular module ${}_R R$.

Definition 17.13. *Let R be a ring. The Jacobson radical of R is the ideal $\text{Rad}({}_R R)$. It is denoted by $J(R)$.*

By the Remarks 17.5 and 17.12, the Jacobson radical of R can be described as the intersection of the annihilators of the simple left R -modules $\text{Ann}_R(S)$. In particular it is two-sided ideal of R .

Lemma 17.14. *For every $M \in R\text{-Mod}$, $J(R)M \leq \text{Rad}(M)$*

Proof. Since $J(R)$ annihilates any simple module S , all homomorphisms $M \rightarrow S$ are zero on $J(R)M$ so, by Remark 17.12, $J(R)M \leq \text{Rad}(M)$ \square

Proposition 17.15 (Nakayma's Lemma). *Let M be a finitely generated R -module. If L is a submodule of M such that $L + J(R)M = M$, then $L = M$.*

Proof. $L + J(R)M = M$ implies $L + \text{Rad}(M) = M$ and since $\text{Rad}(M)$ is superfluous in M (see Proposition ??) we get $L = M$. \square

We conclude with the following characterization of $J(R)$

Proposition 17.16. $J(R) = \{r \in R \mid 1 - xr \text{ has a left inverse for any } x \in R\}$

18. EXERCISE

Exercise 18.1. Let $M \in R\text{-Mod}$ be finitely generated. Show that, for any $L < M$, there exists a maximal submodule of M containing L . In particular, $\text{Rad}(M) < M$.

Exercise 18.2. Show that, for any simple module $S \in R\text{-Mod}$, $\text{Ann}_R(S)$ is a two-sided ideal of R .

Exercise 18.3. Let $p \in \mathbb{N}$ a prime and $M = \{\frac{a}{p^n} \in \mathbb{Q} \mid a \in \mathbb{Z}, n \in \mathbb{N}\}$.

- (1) Verify that $\mathbb{Z} \leq M \leq \mathbb{Q}$ in $\mathbb{Z}\text{-Mod}$.
- (2) Let $\mathbb{Z}_{p^\infty} = M/\mathbb{Z}$. Show that \mathbb{Z}_{p^∞} is a divisible group.
- (3) show that any $L \leq \mathbb{Z}_{p^\infty}$ is cyclic, generated by an element $\frac{1}{p^l}$, $l \in \mathbb{N}$.

Conclude the the lattice of the subgroups of \mathbb{Z}_{p^∞} is a well-ordered chain and so \mathbb{Z}_{p^∞} does not have any maximal subgroup.

19. LOCAL RINGS

Definition 19.1. A ring R is a local ring if all the non-invertible elements form a proper ideal of R .

In other words, setting $U(R) = \{x \in R \mid x \text{ is invertible}\}$, R is a local ring if $R \setminus U(R)$ is a left ideal of R . One easily shows that $R \setminus U(R)$ is a left ideal if and only if it is a two-sided ideal of R (Verify!).

Proposition 19.2. Let R be a local ring. Then

- (1) $R \setminus U(R)$ is the Jacobson radical $J(R)$ of R .
- (2) $R/J(R)$ is a division ring.
- (3) there is a unique simple module (modulo isomorphisms) in $R\text{-Mod}$, $S = R/J(R)$. In particular $E(R/J(R))$ is the minimal injective cogenerator of $R\text{-Mod}$.
- (4) The unique idempotent elements in R are 0 and 1.

Proof. 1) Given a ring R , any left ideal of R is contained in $R \setminus U(R)$. So, if R is local, $R \setminus U(R)$ is the unique maximal ideal of R . In particular $R \setminus U(R)$ is the Jacobson radical $J(R)$ of R .

2) is obvious, since every element in $R/J(R)$ is invertible.

3) It follows since $J(R)$ is the unique maximal ideal of R .

4) Let e an idempotent element in a ring R . Observe that from $e(1-e) = 0$, if e is invertible one gets $e = 1$. So, if R is local and e is a not invertible idempotent, then $e \in R \setminus U(R) = J(R)$ and so the idempotent $1-e \in U(R)$ (otherwise we would have $1 \in J(R)$). Hence, $1-e = 1$ and so $e = 0$. We conclude that the only idempotents in R are the trivial ones, i.e. 0 and 1. \square

Remark 19.3. If R is a local ring, then ${}_R R$ is an indecomposable R -module, since the direct summands of ${}_R R$ correspond to the idempotent elements of R (see Exercise 9.4).

If $M \in R\text{-Mod}$ and $\text{End}_R(M)$ is a local ring, then M is indecomposable. Indeed, to any decomposition $M = N \oplus L$, we can associate an idempotent element $\pi_N \in \text{End}_R(M)$, $\pi_N : M \rightarrow M$, $n + l \mapsto n$. Thus $\pi_N = 0$ or $\pi_N = \text{id}_M$ in $\text{End}_R(M)$, from which we get $N = 0$ or $N = M$, respectively.

20. FINITE LENGTH MODULES

Let $M \in R\text{-Mod}$. A sequence $0 = N_0 \leq N_1 \leq \dots \leq N_{s-1} \leq N_s = M$ of submodules of M is called a *filtration* of M , with *factors* N_i/N_{i-1} , $i = 1, \dots, s$. The *length* of the filtration is the number of non-zero factors.

Consider now a filtration $0 = N'_0 \leq N'_1 \leq \dots \leq N'_t = M$; it is a *refinement* of the latter one if $\{N_i \mid 0 \leq i \leq s\} \subseteq \{N'_i \mid 0 \leq i \leq t\}$.

Two filtrations of M are said *equivalent* if $s = t$ and there exists a permutation $\sigma : \{0, 1, \dots, s\} \rightarrow \{0, 1, \dots, s\}$ such that $N_i/N_{i-1} \cong N'_{\sigma(i)}/N'_{\sigma(i)-1}$, for $i = 1, \dots, s$.

Finally, a filtration $0 = N_0 \leq N_1 \leq \dots \leq N_{s-1} \leq N_s = M$ of M is a *composition series* of M if the factors N_i/N_{i-1} , $i = 1, \dots, s$, are simple modules. In such a case they are called *composition factors* of M .

Theorem 20.1. Any two filtrations of M admit equivalent refinements.

Proof. The proof follows from the following Lemma: *Let $U_1 \leq U_2 \leq M$ and $V_1 \leq V_2 \leq M$. Then $(U_1 + U_2 \cap V_2)/(U_1 + V_1 \cap U_2) \cong (U_2 \cap V_2)/(U_1 \cap V_2) + (U_2 + V_1) \cong (V_1 + U_2 \cap V_2)/(V_1 + U_1 \cap V_2)$*

In our setting, consider $0 = N_0 \leq N_1 \leq \dots \leq N_{s-1} \leq N_s = M$ and $0 = L_0 \leq L_1 \leq \dots \leq L_{s-1} \leq L_t = M$ two filtrations of M . For any $1 \leq i \leq s$ and $1 \leq j \leq t$ define $N_{i,j} = N_{i-1} + (L_j \cap N_i)$ and $L_{j,i} = L_{j-1} + (N_j \cap L_i)$. Then

$$0 = N_{1,0} \leq N_{1,1} \leq \dots \leq N_{1,t} \leq N_{2,0} \leq \dots \leq N_{2,t} \leq \dots \leq N_{s,t} = M$$

is a refinement of the first filtration with factors $F_{i,j} = N_{i,j}/N_{i,j-1}$ and

$$0 = L_{1,0} \leq L_{1,1} \leq \dots \leq L_{1,s} \leq L_{2,0} \leq \dots \leq L_{2,s} \leq \dots \leq L_{t,s} = M$$

is a refinement of the second filtration with factors $G_{j,i} = L_{j,i}/L_{j,i-1}$. Clearly the two refinements have the same length st and by the stated lemma $F_{i,j} \cong G_{j,i}$. \square

As a corollary of the previous Theorem, we get the following crucial result, known as Jordan-Hölder Theorem:

Theorem 20.2 (Jordan-Hölder). *Let $M \in R\text{-Mod}$ a module with a composition series of length l . Then*

- (1) *Any filtration of M has length at most l and it can be refined in a composition series of M .*
- (2) *All the composition series of M are equivalent and have length l .*

Proof. The proof follows by the previous proposition, since a composition series does not admit any non trivial refinement. \square

This leads to the following definition:

Definition 20.3. *A module $M \in R\text{-Mod}$ is of finite length if it admits a composition series. The length l of any composition series of M is called the length of M , denoted by $l(M)$.*

- Example 20.4.*
- (1) Any vector space of finite dimension over a field K is a K -module of finite length. Its length coincides with its dimension.
 - (2) The regular module ${}_Z\mathbb{Z}$ is not of finite length.

In the following proposition we collect some relevant properties of finite length modules: some of them are trivial, some of them need a short proof that we leave for exercise.

Proposition 20.5. *Let $M \in R\text{-Mod}$ be a finite length module. Then*

- (1) *M is finitely generated*
- (2) *for any $N \leq M$, N and M/N are of finite length*
- (3) *If $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ is an exact sequence, then $l(M) = l(N) + l(L)$*
- (4) *M is a direct sums of indecomposable submodules.*
- (5) *$\text{Soc}(M)$ is an essential submodule of M*
- (6) *$M/\text{Rad}(M)$ is semisimple (i.e. direct sum of simple modules)*
- (7) *M contains a finite number of simple modules*

Proof. 4) If M is indecomposable the statement is trivially true. Otherwise we argue by induction on $l(M)$. If $M = V_1 \oplus V_2$, by point 3) we get that $l(V_1) < l(M)$ and $l(V_2) < l(M)$, so V_1 and V_2 are direct sums of indecomposable submodules.

5) Any $L \leq M$ has a composition series, so it contains a simple submodule, which is of course also a simple submodule of M .

6) By induction on $l(M/\text{Rad}(M))$

7) Any simple submodule of M is a direct summand of $\text{Soc}(M)$. Since $\text{Soc}(M)$ is finitely generated (by (1) and (2)), it has only a finite number of summands. \square

For modules of finite length the converse of Remark 19.3 holds.

Lemma 20.6. *Let $M \in R\text{-Mod}$ a module of finite length $l(M) = n$. Then, for any $f : M \rightarrow M$, one has $M = \text{Im } f^n \oplus \text{Ker } f^n$.*

Proof. Consider the sequence of inclusions $\cdots \leq \text{Im } f^2 \leq \text{Im } f \leq M$. Since M has finite length, the inclusions are trivial for almost every $i \in \mathbb{N}$. In particular, there exists m such that $\text{Im } f^m = \text{Im } f^{2m}$ and we can assume $m = n$. Let now $x \in M$: hence $f^n(x) = f^{2n}(y)$ for $y \in M$ and so $x = f^n(y) - (x - f^n(y)) \in \text{Im } f^n + \text{Ker } f^n$.

Moreover, from the sequence of inclusions $0 \leq \text{Ker } f \leq \text{Ker } f^2 \leq \cdots \leq M$, arguing as before we can assume $\text{Ker } f^n = \text{Ker } f^{2n}$. Consider now $x \in \text{Im } f^n \cap \text{Ker } f^n$. So $x = f^n(y)$ and $f^n(x) = f^{2n}(y) = 0$. Hence $y \in \text{Ker } f^n$ and so $x = f^n(y) = 0$. \square

Proposition 20.7. *Let $M \in R\text{-Mod}$ an indecomposable module of finite length. Then $\text{End}_R(M)$ is a local ring*

Proof. Let $f : M \rightarrow M$. Since M is indecomposable, by the previous lemma one easily conclude that f is a mono if and only if it is an epi if and only if it is an iso if and only if $f^m \neq 0$ for any $m \in \mathbb{N}$ (see Exercise 21.1).

Thus let $U = \{f \in \text{End}_R(M) \mid f \text{ is invertible}\}$. Let us show that $\text{End}_R(M) \setminus U$ is an ideal of $\text{End}_R(M)$. So let f, g in $\text{End}_R(M) \setminus U$. The crucial point is to show that $f + g$ is not invertible (see Exercise 21.1). If $f + g$ would be invertible, there would exist $h \in U$ such that $(f + g)h = \text{id}_M$. Since $g \notin U$, then $gh \notin U$, so gh would be nilpotent. Let r such that $(gh)^r = 0$: from $(\text{id}_M - gh)(\text{id}_M + gh + (gh)^2 + \cdots + (gh)^{r-1}) = \text{id}_M$ we would conclude $fh \in U$ and so $f \in U$. \square

Theorem 20.8 (Krull-Remak-Schmidt-Azumaya). *Let $M \cong A_1 \oplus A_2 \oplus \cdots \oplus A_m \cong C_1 \oplus C_2 \oplus \cdots \oplus C_n$ where $\text{End}_R(A_i)$ is a local ring for any $i = 1, \dots, m$ and C_j is indecomposable for any $j = 1, \dots, n$. Then $n = m$ and there exists a bijection $\sigma : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ such that $A_i \cong C_{\sigma(i)}$ for any $i = 1, \dots, n$.*

Proof. By induction on m .

If $m = 1$, then $M \cong A_1$ is indecomposable and so we conclude.

If $m > 1$, consider the equalities

$$\text{id}_{A_m} = \pi_{A_m} i_{A_m} = \pi_{A_m} \left(\sum_{j=1}^n i_{C_j} \pi_{C_j} \right) i_{A_m} = \sum_{j=1}^n \pi_{A_m} i_{C_j} \pi_{C_j} i_{A_m},$$

where π and i are the canonical projections and inclusions. Since $\text{End}_R(A_m)$ is local, and in any local ring the sum of not invertible elements is not invertible, there exist \bar{j} such that $\alpha = \pi_{A_m} i_{C_{\bar{j}}} \pi_{C_{\bar{j}}} i_{A_m}$ is invertible. We can assume $\bar{j} = n$, and consider $\gamma = \alpha^{-1} \pi_{A_m} i_{C_n} : C_n \rightarrow A_m$. Since $\gamma \pi_{C_n} i_{A_m} = \alpha^{-1}$, we get that γ is a split epimorphism. Since C_n is indecomposable, we conclude γ is an iso, and so $C_n \cong A_m$. Then apply induction to get the thesis. \square

The previous theorem says that if M is a module which is a direct sum of modules with local endomorphism rings, then any two direct sum decompositions of M into indecomposable direct summands are isomorphic. We conclude that the modules of finite length admits a unique (modulo isomorphisms) decomposition in indecomposable modules

21. EXERCISES

Exercise 21.1. Let M an indecomposable R -module of finite length and $f \in \text{End}_R(M)$. Show that the following are equivalent:

- (1) f is a mono
- (2) f is an epi
- (3) f is an iso
- (4) f is not nilpotent.

In particular, if f is not invertible, then gf is not invertible for any $g \in \text{End}_R(M)$. Which of the previous implications hold also if M is of finite length but not indecomposable?

Exercise 21.2. Let M be an R -module.

- (1) Let $M_1, M_2 \leq M$ such that $M_1 + M_2 = M$. Show that $M/M_1 \cap M_2 \cong M_1/M_1 \cap M_2 \oplus M_2/M_1 \cap M_2$.
- (2) Suppose $\text{Rad}(M) = M_1 \cap M_2$, where M_1 and M_2 are maximal submodules of M . Show that $M/\text{Rad}(M) = S_1 \oplus S_2$ where S_1 and S_2 are simple R -modules.
- (3) Let M be a finite length R -module. Show that $M/\text{Rad}(M)$ is semisimple.

22. FINITE DIMENSIONAL K -ALGEBRAS

Definition 22.1. Let K be a field. A K -algebra Λ is a ring with a map $K \times \Lambda \rightarrow \Lambda$, $k \mapsto ka$, such that Λ is a K -module and $k(ab) = a(kb) = (ab)k$ for any $k \in K$ and $a, b \in \Lambda$. Λ is finite dimensional if $\dim_K(\Lambda) < \infty$.

In other words, a K -algebra is a ring with a further structure of K -vector space, compatible with the ring structure.

Remark 22.2. Any element $k \in K$ can be identify with an element of Λ by means of $K \times \Lambda \rightarrow \Lambda$, $k \mapsto k \cdot 1$. Thanks to this identification, we get that $K \leq \Lambda$ so any Λ -module is in particular a K -module.

Example 22.3. (1) The ring $M_n(K)$ is a finite dimensional K -algebra. with $\dim_K(M_n(K)) = n^2$. Any element $k \in K$ is identified with the diagonal matrix with k on the diagonal elements.

(2) The ring $K[x]$ is a K -algebra, not finite dimensional.

Proposition 22.4. Let Λ be a finite dimensional K -algebra. Then $M \in \Lambda\text{-Mod}$ is finitely generated if and only if $\dim_K(M) < \infty$.

Proof. Assume $\dim_K(\Lambda) = n$ and $\{a_1, \dots, a_n\}$ a K -basis.

If $\{m_1, \dots, m_r\}$ is a set of generator of M as Λ -module, then one verifies that $\{a_i m_j\}_{i=1, \dots, n}^{j=1, \dots, r}$ is a set of generators for M as K -module.

Conversely, if M is generated by $\{m_1, \dots, m_s\}$ as K -module, since $K \leq \Lambda$, one gets that M is generated by $\{m_1, \dots, m_s\}$ also as Λ -module. □

In the following we denote by $\Lambda\text{-mod}$ the full subcategory of $\Lambda\text{-Mod}$ consisting on the finitely generated Λ -modules.

Corollary 22.5. Any finitely generated module $M \in \Lambda\text{-mod}$ is a finite length module, and $l(M) \leq \dim_K(M)$.

Proof. Since any $M \in \Lambda\text{-mod}$ is a finite dimensional vector space, M admits a composition series in $K\text{-mod}$ of length n , where $\dim_K(M) = n$. So any filtration of M in $\Lambda\text{-Mod}$ is at most of length n and any refinement is a refinement also in $K\text{-mod}$. Thus we conclude. □

Proposition 22.6. Let $M, N \in \Lambda\text{-mod}$. Then $\text{Hom}_\Lambda(M, N)$ is a finitely generated K -module. In particular, $\Gamma = \text{End}_\Lambda(M)$ is a finite dimensional K -algebra and M_Γ is finitely generated.

Proof. The K -module $\text{Hom}_\Lambda(M, N)$ is a K -submodule of $\text{Hom}_K(M, N)$, and the latter is finitely generated by a well-known result of linear algebra. Thus $\text{Hom}_\Lambda(M, N)$ is finitely generated as K -module. In particular, $\Gamma = \text{Hom}_\Lambda(M, M)$ is a finite dimensional K -algebra. Since M has a natural structure of right Γ -module and it is a finitely generated K -module, it is also a finitely generated Γ -module. □

In the sequel, let Λ be a finite dimensional K -algebra.

Since ${}_\Lambda\Lambda$ is of finite length, it admits a unique decomposition in indecomposable submodules. The indecomposable summands of a ring are in correspondence with the idempotent elements, so there exists a set $\{e_1, e_2, \dots, e_n\}$ of idempotents of Λ such that ${}_\Lambda\Lambda = \Lambda e_1 \oplus \dots \oplus \Lambda e_n$. Moreover we can assume $1 = e_1 + \dots + e_n$ and one easily shows that $e_i e_j = 0$ for any $i \neq j$ (a set of idempotents with this property is called *orthogonal*). Finally since Λe_i are indecomposable, each idempotent e_i is *primitive* (i.e. it cannot be a sum of two non-zero orthogonal idempotents, see Exercise 23.1). Notice that ${}_\Lambda\Lambda = e_1 \Lambda \oplus \dots \oplus e_n \Lambda$ is a decomposition in indecomposable summands of the regular right module ${}_\Lambda\Lambda$. From this discussion it follows that, for $i = 1, \dots, n$, the $P_i = \Lambda e_i$ are the indecomposable projective left Λ -modules and the $Q_i = e_i \Lambda$ are the indecomposable projective right Λ -modules (Why?).

Consider the functor $D : \Lambda\text{-mod} \rightarrow \text{mod-}\Lambda$, $M \mapsto D(M) = \text{Hom}_K({}_\Lambda M, K)$. For simplicity, we denote by D the analogous functor $D : \text{mod-}\Lambda \rightarrow \Lambda\text{-mod}$, $N \mapsto D(N) = \text{Hom}_K(N_\Lambda, K)$. For any $M \in \Lambda\text{-mod}$ define the *evaluation morphism* $\delta_M : M \rightarrow D^2(M)$, $x \mapsto \delta_M(x)$, where $\delta_M(x) : D(M) \rightarrow K$, $\varphi \mapsto \varphi(x)$. One easily verify that δ_M is an isomorphism for any $M \in \Lambda\text{-mod}$. Similarly one define δ_N for any $N \in \text{mod-}\Lambda$, which is an iso for any N .

It turns out that $\delta : D^2 \rightarrow 1$ is a natural transformation which defines a duality between $\Lambda\text{-mod}$ and $\text{mod-}\Lambda$. So P is indecomposable projective in $\Lambda\text{-mod}$ if and only if $D(P)$ is indecomposable injective in $\text{mod-}\Lambda$. S is simple in $\Lambda\text{-mod}$ if and only if $D(S)$ is simple in $\text{mod-}\Lambda$ (Why?)

Thanks to the duality (D, D) , we conclude that $D(\Lambda_\Lambda)$ is the minimal injective cogenerator of $\Lambda\text{-mod}$ and the $E_i = D(Q_i)$ are the unique indecomposable injective modules in $\Lambda\text{-mod}$. Observe that if S and T are non isomorphic simple modules in $\Lambda\text{-mod}$, then their injective envelopes $E(S)$ and $E(T)$ are non isomorphic indecomposable injective modules (Why?). We conclude that there is a finite number of simple left Λ -modules S_1, S_2, \dots, S_n .

Observe that in $\text{mod-}\Lambda$ there exist injective envelopes so, thanks to the duality, we get that in $\Lambda\text{-mod}$ there exist projective covers. Let us see how to compute injective envelopes and projective covers.

First observe that, denoted by $J = J(\Lambda)$, for any $M \in \Lambda\text{-mod}$ the submodule JM is superfluous in M 17.11. In particular Je_1 is superfluous in Λe_i , so Λe_i is the projective cover of $\Lambda e_i / Je_i$ (see 12.8). Moreover, since Λe_i is indecomposable, we get that $\Lambda e_i / Je_i$ is a simple module (see Exercise 23.2) and so Je_1 is a maximal submodule of Λe_i . We conclude that $S_i = \Lambda e_i / Je_i$ is a complete list of the simple modules in $\Lambda\text{-mod}$. Similarly, $T_i = e_i \Lambda / e_i J$ is a complete list of the simple modules in $\text{mod-}\Lambda$. Notice that, as a consequence of the the above discussion, we get that Je_1 is the radical of Λe_i (Why?). One can show that the same result holds for any $M \in \Lambda\text{-mod}$: $\text{Rad}(M) = JM$.

Since $S_i = D(T_i)$, we get that $E_i = D(Q_i)$ is the injective envelope of S_i .

How to compute injective envelopes and projective covers for any $M \in \Lambda\text{-mod}$? Since it is of finite length, $M/\text{Rad}(M)$ and $\text{Soc}(M)$ are semisimple. Let $M/\text{Rad}(M) = S_1 \oplus \dots \oplus S_r$ (eventually with a certain multiplicity). Then $P(M) = P(S_1) \oplus \dots \oplus P(S_r)$. Similarly, if $\text{Soc}(M) = S_1 \oplus \dots \oplus S_m$, then $E(M) = E(S_1) \oplus \dots \oplus E(S_m)$. (see Exercises 23.3 and 23.4).

To conclude: in $\Lambda\text{-mod}$ the simples are the $S_i = \Lambda e_i / Je_i$, the indecomposable projectives are the $P_i = \Lambda e_i$, the indecomposable injectives are the $E_i = D(e_i \Lambda)$. The regular module ${}_\Lambda \Lambda$ is the minimal projective generator of $\Lambda\text{-mod}$ and $D(\Lambda_\Lambda)$ is the minimal injective cogenerator of $\Lambda\text{-mod}$. Moreover P_i is the projective cover of S_i and E_i is the injective envelope of S_i .

In $\text{mod-}\Lambda$ the simples are the $T_i = \Lambda e_i / Je_i = D(S_i)$, the indecomposable projectives are the $Q_i = e_i \Lambda$, the indecomposable injectives are the $F_i = D(\Lambda e_i)$. The regular module Λ_Λ is the minimal projective generator of $\text{mod-}\Lambda$ and $D(\Lambda_\Lambda)$ is the minimal injective cogenerator of $\text{mod-}\Lambda$. Moreover Q_i is the projective cover of T_i and F_i is the injective envelope of T_i .

23. EXERCISES

Exercise 23.1. A idempotent element $e \in \Lambda$ is called *primitive* if it is not a sum of two non zero orthogonal idempotents. Show that Λe is indecomposable if and only if e is primitive.

Exercise 23.2. Let Λ a finite dimensional algebra. Let $M = N_1 \oplus N_2$ and assume that P_1 and P_2 are projective covers of N_1 and N_2 , respectively. Show that $P_1 \oplus P_2$ is the projective cover of M . Similarly, assume that E_1 and E_2 are the injective envelopes of N_1 and N_2 , respectively, then $E_1 \oplus E_2$ is the injective envelope of M .

Exercise 23.3. Let $M \in \Lambda\text{-mod}$ and $\text{Soc}(M) = S_1 \oplus \dots \oplus S_r$. Show that there exists an essential monomorphism $0 \rightarrow M \rightarrow E(S_1) \oplus \dots \oplus E(S_r)$ and conclude that $E(M) = E(\text{Soc}(M)) = E(S_1) \oplus \dots \oplus E(S_r)$. (Hint: $\text{Soc}(M)$ is essential in M , so...)

Exercise 23.4. Let $M \in \Lambda\text{-mod}$ and $M/\text{Rad}(M) = S_1 \oplus \dots \oplus S_r$. Show that there exists a superfluous epimorphism $M \rightarrow P(S_1) \oplus \dots \oplus P(S_r) \rightarrow 0$ and conclude that $P(M) = P(M/\text{Rad}(M)) = P(S_1) \oplus \dots \oplus P(S_r)$. (Hint: $\text{Rad}(M)$ is superfluous in M , so...)